

RECORD

F. R. Fosberg

Collection and Field Note Book

Page	Description
1-4	Florida
1-4	Hwy 1000
4-5	Specimens collected
6-11	Flight Homestead AFB to Dry Tortugas
12-14	Tortugas
15-19	Specimens collected
19-21	Bush Key
21-23	Garden Key
23-25	Loganhead Key
25-27	Long Key
27-30	Garden Key
30-31	South of Homestead
31-32	Key Largo
32-33	Trip Miami to Key Largo
33-34	North of Jewish Cr.
34-35	District of Columbia-Kentworth Gann
35	Virginia
35-36	Grimey Springs
36-37	Shenandoah Nat. Pk.
37-38	Old Rag View
38-39	Big Meadows
39-40	...
40-41	...
41-42	...
42-43	...
43-44	...
44-45	...
45-46	...
46-47	...
47-48	...
48-49	...
49-50	...
50-51	...
51-52	...
52-53	...
53-54	...
54-55	...
55-56	...
56-57	...
57-58	...
58-59	...
59-60	...
60-61	...
61-62	...
62-63	...
63-64	...
64-65	...
65-66	...
66-67	...
67-68	...
68-69	...
69-70	...
70-71	...
71-72	...
72-73	...
73-74	...
74-75	...
75-76	...
76-77	...
77-78	...
78-79	...
79-80	...
80-81	...
81-82	...
82-83	...
83-84	...
84-85	...
85-86	...
86-87	...
87-88	...
88-89	...
89-90	...
90-91	...
91-92	...
92-93	...
93-94	...
94-95	...
95-96	...
96-97	...
97-98	...
98-99	...
99-100	...

Collection and Field Note Book

No. 65

(Aug. 1, 1962 - Oct. 10, 1962)

(42973 - 43244)


	<u>Pages</u>
<u>Florida</u>	<u>1-47</u>
Everglades Nat'l Park.....	1-4
Homestead Air Force Base	
Specimen collected..42973	4-5
Flight Homestead AFB to Dry Tortugas	6-11
Tortugas.....	12-41
Specimens collected	
Bush Key	42974-42978
	16-19
	42998-43000
	18-21
Garden Key ...	42979-42997
	16-19
	43008-43025
	20-23
 24-25, 36
Loggerhead Key	26-30
	43026-43053.....
	32-35
Long Key	43001-43007.....
	20-21
Garden Key... 43054-43069	38-40
south of Homestead	
	43070-43072.....
	40-41
Key Largo.....	43073
	40-41
	43074-43083
	40-41, 46-47
Trip Miami to Key Largo	42-44
North of Jewfish Cr. 43084-43086..	46-47
District of Columbia-Kenilworth Gdns	50-51

Virginia

Orkney Springs	52
Shenandoah Nat. Pk	53-65
<u>Old Rag View...</u> 43091-43098	54-55
Big Meadows... 43099-43102	54-57
	43103-43111
	56-57
	43112-43129
	60-63
 58-59
Swift Run Gap 43130-43138	62-65
Near Williamsburg 43232-43235	128-129
(Powhatan Creek-swamp)	

Virginia (continued)

<u>Shenandoah National Park</u>	
Pinnacles Picnic Ground..43139	64-65
<u>Flight Minneapolis to Portland</u>	
<u>Minnesota</u> (from air)	66
Dakotas... (from air)	67
Montana....(from air)	68-73
Idaho (from air)	74-76
Oregon ... (from air)	77
Washington (from air)	77-79
Ohio...Drysart Woods,. 43237	130-131, 134
<u>Oregon -- trip from Corvallis to coast...</u>	80-83
North of Coquille ... 43218	116
Toledo	43140-43155
	84-87
Yaquina Bay Bridge ...43156-43170	86-89
Waldport	43171-43182
	88-91
Florence-Mercer Lk Rd 43183-43188	90-91
(Lincoln County)	
Triangle Lake	43189-43200
	96-97
Cape Perpetua State Park.....	92
Hauser(Gees.County) 43201-43217	114-117
Trip Corvallis-Arcata via Eugene,	
Reedsport.....	98-102
California	103-121
Del Norte Redwood Grove	103
Auto trip Eureka to Bull Creek	104-107
Air trip Arcata-San Francisco	108-113
Hayward (Alameda Cty) 43219.....	116-117
Santa Cruz Mts. Ridge 43220-43231...	116-119
Flight San Francisco to Chicago.....	120-127
via Nevada, Utah	
<u>Virginia</u>	
Sleepy Hollow.... 43236, 43087-43090	130-131, 48-49
Falls Church area	136
Rattlesnake Point Overlook.....	138
	43243.....
	138-139
Washington, D. C.... 43244	138-139
<u>Maryland -- Arnold..43238-43242-----</u>	130-131, 137
Suitland Bog....	135
Broad Creek(Anne Arundel Co.)	136



VANCOUVER
Pt. Grey
University Hill
English B.

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

WESTMINSTER
Richmond
Surrey
Langley

FRASER MILLS
Maple Ridge
Mission
Matsqui

CHILLIWACK
Smithvale
Rosedale
Sardis

CHILLIWACK
Vedder
Cultus Lake
Hurry Park

CHILLIWACK
Sumas
Abbotsford
Huntingdon

CHILLIWACK
Sumas
Hilltop
Columbia

CHILLIWACK
Nooksack
Kendall
Lawrence

CHILLIWACK
Bellingham
Geneva
Tawanda

CHILLIWACK
Sumas
Hilltop
Columbia

ITINERARY ----- COAST FIELD TRIP

FRIDAY, AUGUST 24 (All Times Daylight Saving)

- 8:00 Assemble, 30th Street and Campus Way, O. S. U. campus. Buy lunches, if desired--\$1.00. Riders will be assigned to cars having space.
- 8:30 Leave O. S. U. campus, proceed west on Highway 20, via Philomath, Blodgett, Burnt Woods, Eddyville, to Toledo. At Toledo, turn left (south) on old road to Newport.
- 10:15 Arrive at first stop, at salt marsh between Toledo and Newport. Park in single line along shoulder of road.
- 11:30 Leave salt marsh. Proceed through Newport to Yaquina Bay (Newport) State Park, just northwest of Yaquina Bay Bridge, for lunch.
- 1:00 Leave Newport, go south on Highway 101 to a point ca. 2 miles south of Yaquina Bay Bridge. There we will park in single line along shoulder of highway and walk west through some woods to an area of marsh and sandy beach.
- 2:30 Leave this locality, go south on Highway 101 ca. 13 miles to a spot ca. 1 mile north of Waldport. Park in a line along shoulder of highway, and walk west toward ocean to see coastal forest and dunes.
- 4:00 Leave here and proceed south, through Waldport, toward Yachats. Check in at motel (The Adobe or Deane's OceanSide, both a short distance north of Yachats) or find camping site at one of 3 suggested state parks near Yachats.

SATURDAY, AUGUST 25

- 8:00 Assemble at The Adobe; use same travel arrangements as on Friday, unless otherwise agreed upon.
- 8:15 Leave The Adobe, proceed south through Yachats to Cape Perpetua State Park. We will go first up the high road to the lookout point, then down and into main part of park. Drive to parking circle at end of the road.
- 10:15 Leave Cape Perpetua Park, drive south on Highway 101 ca. 19 miles to the Mercer Lake turnoff, 5 miles north of Florence. Turn left (east) and go for ca. 1/4 mile; park in single line on shoulder of road. We will walk down to a Darlingtonia bog, through a forest of pines, cedars, and hemlocks.
- 12:00 Leave the bog and drive south to Florence. Lunch can be purchased at various restaurants there. We will use Goddard's Restaurant, 2 miles south of Florence, as a rallying point.
- 1:00 Meet at Goddard's Restaurant to proceed south from Florence 8 miles to Carter Lake Forest Camp. This is an area of active dunes, encroaching on a coastal forest.
- 2:30 Leave Carter Lake and return north to Florence on Highway 101. The shortest return to Corvallis is east via Highway 36 (shown on your mimeographed map as U.S. 28). It is 90 miles from Florence to Corvallis, and a good two hours' drive. Highway 36 joins Highway 99W, north of Eugene, and you can follow the latter highway north to Corvallis; or, if you are alert, there is a connecting road north to Corvallis at Cheshire (on Highway 36, 4 miles before the junction with 99W); Cheshire is ca. 63 miles from Florence.

On Highway 36 there are some possible stopping points not on our formal itinerary. From 1 to 3 miles past Swisshome, 24 to 27 miles from Florence, there are some high, wet cliffs along the road that could be searched for ferns. Farther on, just west of Triangle Lake, ca. 40 miles from Florence, there is a large parking area that gives access to a douglas fir forest.

A car will lead the group back to the University, without making these stops; but if any cars wish to stop, they should have no difficulty finding their way "home" to Corvallis.

Notes:

1. Transportation. The total mileage is approximately 225 miles. Guest riders should make appropriate arrangements to share in travel expenses. A suggested way to handle this is to pay 1¢ per mile per passenger; but some other system might equally well be agreed on.
2. Oregon Highways. These are generally good. However, there are often dangerous curves; and logging trucks are an ever-present menace. I am especially concerned that everyone be most careful getting in and out of cars where we park. We are taking a chance parking on rather narrow shoulders, so one must be alert for oncoming vehicles. At any of the stops on Highway 101, please be sure your car is completely off the pavement when parked.
3. Botanizing. At each stop, we hope you will use the mimeographed list of species to help identify the plants you see. Two of the stops (Cape Perpetua and Carter Lake) are in State Parks, where collecting is not allowed. At other places, I think modest collecting will do no damage. We have been asked to preserve the Darlingtonia plants, however, and not to pick any at the place we stop. There is a spot nearby where someone has bulldozed part of the bog and where some plants of this species may be available (if they haven't already died). Oregon, too, suffers from the virus of creeping commercialization of her shoreline. O. S. U. staff members who will be on the trip are Dr. K. L. Chambers, Dr. William Chilcote, and Miss La Rea Dennis, all of whom will be glad to try to answer questions.
4. Motel. People who have sent money for reservations will be staying at The Adobe. Dr. Chambers will give you your room number and roommate's name. He will contact each of you at the motel Friday evening and, if necessary, either refund part of your payment or collect from you. We hope you will enjoy the ocean scenery from the motel, and will take the chance to botanize on the sea bluffs and examine the rocky tide pools, surge channels, algae, etc.
5. Camping. There are three state parks near Yachats having a total of 102 overnight camp sites. Beachside State Park, 5 miles north of Yachats, has 40; Cape Perpetua, 3 miles south of Yachats, has 48; and Neptune, a mile farther south, has 14. Food can be purchased in Waldport or Yachats.
6. Meals. Box lunches for Friday noon can be bought for \$1.00 each, when we assemble at the O. S. U. campus. There will not be enough for everybody, however, as I expect some families will have their own food with them. There are also restaurants in Newport.
Dinner and breakfast can be obtained at The Adobe or Deane's OceanSide---as well as at other restaurants. Lunch on Saturday should be gotten at restaurants in Florence; we will rally at Goddard's Restaurant, but they can serve only about 30 at a time. Saturday evening after we get back to Corvallis, the O. S. U. Commons Cafeteria will be open to serve us from 5:00 p.m. to 6:00 p.m., or a bit later.

Kenton L. Chambers

LIST OF PLANT SPECIES

TOLEDO SALT MARSH (and vicinity)

Polystichum munitum -- Western Sword-fern
 Pteridium aquilinum -- Western Braken-fern
 Zostera marina -- Eel-grass
 Triglochin maritima -- Seaside Arrow-grass
 Distichlis spicata -- Saltgrass
 Agrostis palustris -- Creeping Bent-grass
 Hordeum brachyantherum -- Meadow Barley
 Anthoxanthum odoratum -- Sweet Vernal-grass
 Deschampsia caespitosa -- Tufted Hair-grass
 Carex sperta & C. hindsii -- Sedge
 Scirpus robustus -- Marsh Bulrush
 Scirpus americanus -- Three-square
 Scirpus microcarpus -- Bulrush
 Juncus effusus -- Common Rush
 Atriplex hastata -- Halberd-leaved Orache
 Salicornia ambigua -- Woody Glasswort
 Potentilla pacifica -- Silver-weed
 Rubus parviflorus -- Thimble Berry
 Oenanthe sarmentosa -- Water Parsley
 Heracleum lanatum -- Cow Parsnip
 Vaccinium ovatum -- Evergreen Huckleberry
 Cuscuta salina var. major -- Salt-marsh Dodder
 Grindelia stricta -- Oregon Gum-plant
 Alnus oregona -- Red Alder
 Rubus laciniatus -- Evergreen Blackberry
 Achillea millefolium var. californica -- Yarrow
 Cotula coronopifolia -- Brass Buttons
 Erechites prenantheoides -- Australian Fireweed
 Epilobium angustifolium -- Fire-weed

Atriplex patula -- Spreading Orache
 Conioselinum chinense -- Hemlock-Parsley
 Orthocarpus castillejoideus -- Owls-clover

NEWPORT MARSH AND SANDY BEACH (and vicinity)

Struthiopteris spicant -- Deer-fern
 Pteridium aquilinum
 Pinus contorta -- Beach pine
 Picea sitchensis -- Sitka spruce
 Poa macrantha -- Seashore Blue-grass
 Elymus mollis -- Sea Lyme-grass
 Calamagrostis nutkaensis -- Pacific Reed-grass
 Agrostis palustris -- Creeping Bent-grass
 Ammophila arenaria -- Beach-grass
 Anthoxanthum odoratum -- Sweet Vernal-grass
 Juncus balticus -- Rush
 Juncus lesueurii -- Salt Rush
 Carex obnupta -- Slough Sedge
 Maianthemum bifolium
 Cardionema ramosissima -- Sand Mat
 Cerastium arvense -- Field Chickweed
 Myrica californica -- Western Wax Myrtle
 Polygonum paronychia -- Beach Knotweed
 Fragaria chiloensis -- Coast Strawberry
 Lathyrus japonicus -- Beach Pea

NEWPORT MARSH -- continued

Potentilla pacifica -- Silver-weed
 Comarum palustre -- Marsh Cinquefoil
 Salix hookeriana -- Willow
 Sisyrinchium californicum
 Abronia latifolia -- Sand-verbena
 Lupinus littoralis -- Seashore Lupine
 Lathyrus littoralis -- Beach pea
 Cakile edentula -- Sea Rocket
 Armeria maritima -- Sea Thrift
 Hypericum anagalloides
 Spiraea douglasii -- Western Spiraea
 Gentiana sceptrum -- Staff Gentian
 Convolvulus soldanella -- Morning-glory
 Centaurium umbellatum -- Centaury
 Plantago lanceolata -- Plantain
 Lonicera involucrata -- Twin-berry
 Rhododendron macrophyllum
 Gaultheria shallon -- Salal
 Arctostaphylos uva-ursi -- Kinnikinnick
 Franseria chamissonis -- Sand-bur
 Hypochaeris radicata -- Cat's-ear
 Achillea millefolium -- Yarrow
 Aster subspicatus -- Aster
 Solidago spathulata -- Sticky Goldenrod
 Vaccinium ovatum -- Evergreen Huckleberry
 Galium trifidum -- Small Bedstraw
 Vaccinium uliginosum -- Blueberry
 Sieglingia decumbens
 Agrostis pallens -- Seashore Bent-grass

WALDFORT DUNES

Most of the above, as well as:

Botrychium silaifolium -- Leathery Grape-^{fern}
 Carex macrocephala -- Large-headed Sedge
 Spiranthes romanzoffiana -- Ladies Tresses
 Habenaria greenei -- Rein Orchid
 Rumex maritimus -- Seaside Dock
 Ranunculus flammula -- Creeping Buttercup
 Sanicula arctopoides -- Snake-root
 Glehnia leiocarpa -- Beach Silver-top
 Angelica hendersonii -- Angelica
 Baccharia pilularis -- Chaparral Broom

CAPE PERPETUA

Polystichum munitum -- Western Sword-fern
 Athyrium filix-femina -- Lady-fern
 Struthiopteris spicant -- Deer-fern
 Polypodium scolieri -- Leather-leaf
 Polypodium vulgare -- Licorice-fern
 Selaginella oregana -- Selaginella
 Equisetum telmateia -- Giant horsetail
 Picea sitchensis -- Sitka Spruce
 Tsuga heterophylla -- Western Hemlock

CAPE PERPETUA -- continued

Maianthemum bifolium -- False Lily-of-the-Valley
Oxalis oregana -- Oregon Oxalis
Lysichitum americanum -- Skunk Cabbage
Ribes bracteosum -- Stink Currant
Rubus spectabilis -- Salmon Berry
Rubus parviflorus -- Thimble Berry
Alnus oregana -- Red Alder
Montia siberica -- Candy Flower
Menziesia ferruginea -- Rustyleaf
Vaccinium parvifolium -- Red Huckleberry
Vaccinium ovatum -- Evergreen Huckleberry
Gaultheria shallon -- Salal
Stachys sp. -- Hedge Nettle
Sambucus glauca -- Blue Elderberry
Sambucus callicarpa -- Red Elderberry
Petasites speciosa -- Western Coltsfoot
Bellis perennis -- Daisy
Scrophularia oregana -- Figwort

DARLINGTONIA BOG AND FOREST

Struthiopteris spicant
Pteridium aquilinum
Pinus contorta -- Coast pine
Thuja plicata -- Western Red Cedar
Tsuga heterophylla -- Western Hemlock
Pseudotsuga menziesii -- Douglas Fir
Myrica californica -- Western Wax Myrtle
Lysichitum americanum -- Skunk-Cabbage
Darlingtonia californica -- California Pitcher-plant
Drosera rotundifolia -- Sundew
Ledum columbianum -- Labrador Tea
Pterospora andromedea -- Pine Drops
Rhododendron macrophyllum -- Rhododendron
Gaultheria shallon -- Salal
Vaccinium ovatum -- Evergreen Huckleberry
Arctostaphylos columbiana -- Hairy Manzanita

F. R. Fosberg
Field notebooks no. 65
begin with no. 42973
end with 43244



Standard Miniature Blank Book

No. 665 9½ x 6 120 Pages Units
No. 667 9½ x 6 200 Pages Units
No. 668 9½ x 6 300 Pages Units

Made in the Following Rulings

Journals, Day or Cash Books, Double \$ and Cts.

S. E. Ledgers, \$ and Cts.

Double Entry Ledgers

Records with Margin Line

When ordering give Number and Ruling desired

Made in U. S. A.

A Boorum & Pease Product

Aug. 1 - Everglades Nat. Park

Penlands south of

a sparse stand of
Pinus elliottii s. *densa*
 (logged off in middle 40's)
 on pitted limestone,
 burned about 3 years ago.
 Pines up to 20m. tall and
 10-15 cm. dbh. bases
 swollen. The pines are
 actually growing in the
 pits in the limestone.
 Ground water showing
 in pits, 5-6 dm down.

For some distance inland
 from Flamingo most of
 the mangroves have been
 killed almost completely
 by Hurricane Donna, Sept. 1960.
Rhizophora, *Sonneratia*,
Avicennia are over large
 areas almost completely
 dead, though still standing.
 Some living *Conocarpus*,
 battered ~~but~~ and completely
 defoliated in the storm,
 now putting out abundant
 leafy branches - these on
 slightly higher ground, where
Conocarpus naturally grows. In
 swamps now a blanket of *Batis*
 and a few scattered mangrove seedlings.

Capeable prairie west of Flamings, - a flat saline mosaic of *Borrichia*, *Spartina*, *Batis*, *Sporobolus virginicus* open mud flats with salt, - on a dense very fine stiff clay-like marl. Patches of forest - badly hit by hurricanes - conditions change drastically from inundation by fresh water, long dry periods, high salt content after hurricanes - when examined the surface was dusty (dust here said to form to 0.7 m. high) but in lowest spots the marl is moist 3-4 cm down. Dark gray, salty to taste.

Paspalum distichum where old cotton eradication camp stood. Other areas, too.

A very slight beach ridge of similar marl, perhaps only a few cm. difference, has a row of *Conocarpus*.

Surface of ground, where bare, dries into a polygonal pattern, but the polygons, rather than indicating simple shrinkage, seem to indicate later expansion, as the cracks bulge.

Inland the hurricane damage is very marked not tapering off gradually. Then a large area of rather less damage.

A broad zone around the main mangrove swamp has low bushes of *Rhizophora*, in places forming a mosaic with areas of saw grass. Mostly 1-2 m. tall, some patches more. Becomes very sparse inland, scattered very small ones in the cladium, well inland these are mainly seedlings.

Saw grass near edge of pinelands ~~is~~ is on a marl surface just at water table. Scattered cobbles of limestone.

In saw grass are scattered clumps of shrubs and trees - mostly bumps in the limestone surface. Some are peat elevations. Clumps of *Taxodium* are basins.

Large areas of sparse stands of dwarfed *Taxodium* (ground spruce type), up to 3-4 m. tall, 4-7 cm. dbh.

scattered in saw grass
trees are very old, at least
have very thin rings, acc.
Robertson.

The pattern of
zonation of the mangrove
species, acc. Robertson,
is very confused and
not at all clear in the
park area.

The death of the
Rhizophora, Laguncularia,
and Avicennia over large
areas resulting from
the hurricane, is very
remarkable, as in many
places the trees seem not
to be ~~or~~ obviously injured (just
as in Guam).

42973

aug 2-

Homestead Air Force Base

42973

Hedyotis nigricans var.common on compacted ground
along strip.

flowers white or almost so

Aug. 2 - flight from Homestead
A7B to Dry Tortugas.

8:28

left field.

Pine lands around Homestead
are very rapidly being cleared
for tomato culture. Only a
relatively few blocks left.

In edges of everglade areas
are evidences of attempts
at agriculture - parallel patterns,
blocks etc.

In parts a fairly large
area of pine with a few scattered
hammocks of denser vegetation.
The pine is mostly open to
semi-open. (photos)

Two long strings of
woody vegetation in the
everglade saw grass area
somewhat broken (photo) - the
alignment looks almost
as though it were not
a natural feature. Between
patches of it nothing seems
visible except the saw grass.

Beyond Mahogany Hammock
there is more water than
land - the small mangrove
belt is very obvious, but
with islands of apparently
hammock vegetation. Then
a maze of mangrove
and tidal channels and
ponds (photos).

Then a strip of mostly
open water.

The part toward the
Flannings road is solid
mangrove with myriads
of tiny ponds. The coastal
portion perhaps 1/5 the
width of the mangrove
swamp, is light gray -
mangroves killed by
Hurricane Donna.

The water in the larger
water bodies in the swamp
is a blackish dull brown.
In some of the smaller ponds
it is chocolate brown.

The Bates in the destroyed
mangrove is rather local.
Apparently much
Bates and other low vegetation,
perhaps floating, in the
neighborhood of the old
canals. The whole inner
Cape Sable area looks generally
unhealthy. Much snail
in tidal channels in outer
Cape Sable - they look milky
or Cafe au lait color.

Very long sand beach
lines Cape Sable island on
the outer side.

The ocean is a dull pea green
off the Cape. Some distance out
it becomes mottled as though
there were some reef growth. Not clear.

The water is not very clear. Gradually becomes too deep to discern a pattern.

Reefs to left - form a very complicated system, but general pattern seems to be more continuous land on outer side, a broken barrier or inner lagoon + cross reefs + channels between, but very irregular.

Sea green color still evident, no bottom pattern visible.

Course more or less parallel to line of Reefs but much to right.

Apparently a strip of deeper water along the inner side of the chain of Reefs, judging by color of water which there is blue rather than green - but seen only from a distance.

Water very gradually deepens along course, too. An abrupt boundary between the milky sea green color and a darker, but still dull, blue. Perhaps has nothing to do with depth.

9:43

Heading directly toward Marquesa ~~Key~~ Atoll -

Water abruptly shallower shortly before we reach M.A. This shallow water forms a wide strip, rather

disscontinuous, with irregularly shallower and deeper water, muds. On this shallow strip are what appear to be submerged dunes. (photos - box.)

Reefs very complex between there and Marquesa Key.

on western side some open sand with very sparse veg.

Marquesa Key mostly covered by mangrove with broad outer beach ridges with scrub forest. Lagoon very shallow with conspicuous winding channels like tidal channels. (photos)

West of Marquesa, a wide mass of reefs with submerged dunes (photos). This extends far to west. These dunes undulate and anastomose but trend is e.-w. And there are very obvious n.-s. channels and current patterns.

Must be due to alternating n.-s. tidal currents.?

This is a very wide feature - several miles, and extends many miles to west. Should be studied in

relation to origin of sand keys.
 Westward as areas
 of small "dunes" many
 of them almost like barchans,
 convex northward. This
 locally superimposed
 on much larger
 undulating network
 pattern.

The whole pattern fades
 out and disappears in deeper
 water westward, with
 a prolongation to southwest.
 Water is much deeper
 to west, but still not a
 clear blue.

Largassum (?) floating
 east of Tortugas. Some
 of it seems to form a line
 along an abrupt deepening,
 or at least change in color
 of water running n-s, just
 as East Key, Tortugas, comes in
 view. East Key grassy with
 some bushes.

Middle is a crescent shaped
 bar - no veg.

~~the bar~~ ^{hospital key} a bar with a patch
 of bushes and a patch of grass (?)
 Photos of Garden, Beach, Ring
 Key, Loggerhead in distance.

Loggerhead as a considerable
 growth of Casuarina from
 end to end, rather thick on

north end. Many coconuts
 in center and a row
 toward north end.

Apparently boulder ridges
 on east side of Long Key.

Aug. 2 - Tortugas
Long Key complex

Considerable low scrub
of Avicennia and
much Salicornia on
inner side, pebble
ridges with little
vegetation on outer sides.
One or two Luriana bushes,
small one small
Cassipouira 0.4 m. tall.
Mats of Sesuvium
mostly on low spots.
Occasional Atriplex.
& patches of Euphorbia
mesembrianthemifolia on
slightly elevated pebble
bars. The Salicornia is
all on areas occasionally
flooded, pebble bottoms
or sand and pebbles.
Small numbers of
Rhizophora up to 1 m. tall
in water on inner
slope of more distal part
(Photo)

swarming
with
Coenobita

Sesuvium

and
Euphorbia
mesemb.

around the
pond
the peat
is much
deeper.

Bush Key - Inner depression
a dense thicket of
Laguncularia (Photo,
end of 12-3, inside). No
other plants in this
area except ~~some~~ a few
Rhizophora. Ground bare.
Around this are broad
flats and beach ridges.
The flats covered by
Luriana scrub,
and Opuntia in openings,
and locally Sporobolus
virginicus and ~~grass~~
~~grass~~ Beach ridge
with Sesuvium
mat, flowing
down onto beach.
Some relatively low
places in Sesuvium
and Sporobolus, kept
so by footy terns, ^{scattered} Portulaca.
Some patches on flats
have Uniola.

In the Laguncularia
forest is a thin layer
of peat under the litter,
then loose coarse sand
mixed with humus.
In the widest part of this
forest is a fair sized pond.
At the west end ^{of the thicket} are a few
bushes of Conocarpus and one of

a sterile Solanum. This is surrounded by Suriana scumb 2 m tall.

Separated from the main mangrove thicket is another small pond completely covered by Rhizophora, dense about 4-5 m tall.

Garden Key

Outside the walls and moat of the old fort there are two lobes of sand that have been built up since the construction of the fort.

These have a few Casuarina and Conocarpus trees. Otherwise they are partly covered by weeds and Cyperus, partly by native strand vegetation - Alriplex, Uniola, Sclerovola, Sesuvium, Sporobolus virginicus, Euphorbia ~~hirsuta~~ mesembrianthemifolia, Ipomoea pes-caprae, and Opuntia dillenii.

Patch of Agave sisalana. Several young Phoenix dactylifera. Inside the fort, the parade ground is a mowed grass lawn of Cyperus stenotaphum, Eragrostis ^{domingensis} with scattered planted

Coronul,

Suriana
Tournefortia
Coccoloba (a)

tuba and

Trees of Bursera simaruba, Blechnum ellipticum, Phoenix dactylifera, Tamarindus indicus, Pandanus tectorius, Thespesia populnea, Conocarpus, Cordia sebestena, Cocos, Terminalia catappa, Coccoloba uvifera, Delonix ^{Citharexylum} ^{nerium indicum}

A number of ornamentals are planted here - Malvarisum arborea, Catharanthus roseus, Rhoeo spathacea, Aloe vera, Caesalpinia pulcherrima, Pedilanthus tithymaloideus, Codiaeum variegatum

Hymenocallis littoralis, Setaria purpurea. Several weeds are abundant here - Euphorbia blodgettii, Phyllanthus amarus, Euphorbia hirta, Lida acuta, Eragrostis ciliaris, Conyza canadensis, Bidens leucantha, ~~Metamorphium~~ Melanthera nivea, Cerberus paniciflora, Brachiaria, Lippia nodiflora, Capriaria, Lepidium, Quillardia. Weeds seen outside only -

Euphorbia prostrata
~~Portulaca~~ ~~Portulaca~~

On the beach, east up, are Thalassia testudinum, Diplanthera wrightii, Syngonium filiformis

Pritchardia
naucifera

Citrus
aurantifolia

Mouroumillea
glabra?

Preynia
quadrifida

Breynia
virens

Washingtonia
filifera

Sida
virens

Cyrtostegia
grandiflora

Musa ^{sep.}

Portulaca
oleracea

Abrus
precatorius

Aug. 2 - Bush Key
low sand islet

42974

Solanum

2

rare, in edge of mangrove thicket

3

75

Philoxerus vermicularis (L.) B.

very local in open
luriana scrub

4

76

Euphorbia mesembriaethemifolia Jacq.

common locally on low
coarse sand ridge

3

77

Rhizophora mangle L.

a few trees in wet thicket
of *Laguncularia*

3

78

Scaevola plumieri (L.) L.

small patch on sand flat

29

Aug. 2 Garden Key
on flat ^{sandy} sandy ground

1

79

(gen) *Eustoma exaltata* (L.) G. Don

rare in ^{ruins of} old fort

6

80

Euphorbia blodgettii Engelm. ex

common in ruins of old fort

2

81

Digitaria horizontalis Willd.

occasional on wall of old fort

2

82

Cyperus planifolius L.C. Rich.

occasional ~~at~~ back of shore

3

83

Cenchrus

common in cracks in
old concrete pavement

3

84

Cynops canadensis (L.) Cronq.

common in cracks in old
concrete pavement

2

85

Digitaria

occasional, weedy places

less

than 1 m.

sparsely leafy, green
stemmed shrub 1 m.
tall, diffusely branched, sterile.

Br.

creeping, forming
loose mat.

dwarf shrub to 4-5 dm.
tall, somewhat glaucous.

small tree 4 m. tall,
with arching prop roots.

shrub up to 0.3 m. tall,
leaves fleshy; flowers white.

erect, glaucous; flowers
purple.

Hitchc.

stems ascending, arching
at tips.

culms decumbent, ^{spreading}
central crown.

caespitose

spreading tufts.

erect

stems decumbent.

- 42986 *Melanthera nivea* (L.) Small
common in weedy places
- 2 87 *Bidens* ~~leucantha~~ ^{alba}
occasional in weedy places
- 1 88 *Lonchocarpus oleraceus* L.
rare in weedy places
- 3 89 *Euphorbia heterophylla*
common in weedy places
- 3 90 *Cenchrus*
common in weedy places
- 1 91 *Euphorbia prostrata*
in cracks in old concrete pavement
- 3 92 *Conocarpus erectus* f.
rare (with common green form)
- 4 93 *Conocarpus erectus* L.
common
- 2 94 *Euphorbia glomerifera* ?
occasional in weedy places
- 1 95 *Euphorbia blodgettii* Engelm.
rare in weedy places
- 2 96 *Cynodon dactylon* (L.) Pers.
common
- 3 97 *Stachytarpheta jamaicensis*
common in weedy places

Aug. 2 - Bush Key
on low sandy flat

- 3 98 *Civicornis germinans* (L.) L.
local in low wet place
- 3 99 *Cyperus planifolius* R. Br. L.C. Rich.
rare

- much branched herb,
flowers white, heads
hemispherical.
rays white.
- erect, glaucous
- erect, base of bracted
leaves scarlet.
- stems spreading
- prostrate, purple.
- shrub 2 m. tall,
leaves silvery.
- shrub 3 m. tall,
leaves green.
flowers white.
- ex Hitchc. stems ascending.
forming a mat.
- (L.) Vahl depressed much branched
plant, flowers light purple.
- shrub 2.5 m. tall,
leaves white beneath,
flowers white.
caespitose.

43000 Sporobolus virginicus L.
4 abundant locally

Aug. 2 - Long Key complex

43001 Laguncularia racemosa Gaertn.
3 occasional in mangrove swamp, on coral gravel

2 02 Rhizophora mangle L.
rare in mangrove swamp on coral gravel

1 03 Sporobolus per-cubrae (L.) Roth
one seen on coral gravel above beach.

4 04 Atriplex pentandra (Jacq.) Standl.
common on gravel beach

2 05 Casuarina
one small plant on outer beach ridge

2 06 Suriana maritima L.
several plants on gravel beach ridges

5 07 Salicornia
common in low places, flooded at high tide

Aug. 3 - Garden Key
on flat coral sand

1 08 Portulaca oleracea L.
common

2 09 Portulaca ~~terrestris~~ oleracea L.
common

forming a loose sod, panicles erect.

shrub 3 m. tall,

shrub 2.5 m. tall, with prop roots; sterile.

seedling

ridge - much branched bushy herb.

seedling 0.5 m. tall; sterile.

small bush 0.5 m tall.

erect, stems and spikes terete, fleshy.

prostrate, ^{stems red} fleshy; flowers yellow, 11-12 mm across, ^{petals} stamens 20-22, ^{emarginate}

stems ascending, fleshy, green, flowers yellow 12-14 mm across, petals emarginate, stamens 19-24.

- 43010 *Lippia nodiflora* L.
occasional
- 3 11 *Desmanthus*
occasional
- 3 12 *Sida procumbens*
common
- 1 13 ^{ringo}*Lycopodium filiforme* Kütz.
thrown up on beach
- 1 14 *Phyllanthus crenatus* Schum. & Th.
occasional inside fort
- 1 15 *Euphorbia hirta* L.
common inside old fort
- 1 16 *Sida acuta* Burm. f.
occasional inside old fort
- 2 17 *Eragrostis ciliaris* (L.) R. Br.
occasional inside old fort
- 1 18 *Diplanthera wrightii* (Aubl.) Aubl.
thrown up on beach
- 3 19 *Eragrostis domingensis* (Pers.)
common inside old fort
- 2 20 *Lippia nodiflora* L.
occasional inside old fort
- 3 21 *Brachiaria*
abundant locally
- 1 22 *Gaillardia*
rare in old fort
- 1 23 *Capriaria biflora*
rare in old fort
- 1 24 *Heliotropium parviflorum*
rare in old fort
- 1 25 *Lepidium*
rare in old fort

- prostrate, somewhat fleshy.
- prostrate
- prostrate
- leaves terete.
- erect
-
- erect.
- leaves flat.
- Stand. small tufts
- prostrate, flowers white.
- prostrate to ascending,
from stolons.
- rays yellow, reddish
near disk
- flowers white
- completely dead.

Aug. 3 - Garden Key
Sand on top of old walls
has a number of weeds,
also

- north lobe of sand -
- a Largely a *Uniola* flat
 - l with patches of *Suriana*,
 - l l *Iva*, *Spartina*, one
 - r small *Thespesia*, one
 - small mangrove *Rhizophora*
 - and a small seedling
 - in moat,
 - c *Cyperus*
 - la *Sporobolus*
 - c *Ipomoea tuba*
 - c *Melanthera*
 - lc *Sesbania*
 - (Many large plants
 - leaving small ones)
 - c *Cyrtosperma dactylon*
 - c *Stachytarpheta jama.*
 - d *Desmanthus*
 - c *Sida procumbens*
 - c *Euphorbia mesem.*
 - l *Spartina lillensis*
 - l *Leaevola plumieri*
 - lc *Canavalia*
 - la *Hymenocallis*
 - lc *Euphorbia heterophylla*
 - c *Bidens baccantha alba*
 - lc *Atriplex pentandra*

Aug. 3 - Tortugas to Key West
Sea east of Tortugas a dull
slightly greenish blue -
very quiet.

Marquesas has a
very convincing atoll
shape. (photos)

West of Marquesas
the bottom pattern is
of sand and brown patches
that suggest quantities
of algae

Suggestion of a secondary
lagoon or bar on n.w. side
inside of ring. This is very
shallow, seemingly not
connected with "tidal channel"
in bottom beside it.

Another series of channels
in mangroves leads right
into a channel in the
lagoon bottom.

Remarkable regular
patterns in sand patches
on bottom east of Marquesas.

Boca Grande Key has a
lagoon-like pond. mangroves
on north side, grass (?) on south.
Mangroves over east and west
lobes.

Third large one from end also
has a pond. Small ones
all as solid patches of mangroves.
(to p. 37)

Aug. 3 - Loggerhead Key
 Almost nothing growing
 under Casuarina
 Sporobolus forms lines of
 shoots along shipwrecks.

Casuarina forms an
 open irregular wood
 to 20 m. tall.

Between it thickets
 of Cordia sebestena,
 scrub of Suriana
 and Opuntia.

Along east shore a
 low dune ridge, about
 2 m. above l.t., cut
 to a sharp escarpment
 by waves.

A number of beds of
 submerged beachrocks
 along this shore just
 outside the present beach.

A caulescent agave
 with antrose prickles
 on edges of lvs. forms
 thickets in the Suriana
 Agave sisalana, also
 forms patches.

Suriana scrub over
 most of south end
 which is open, free from

ms.
 about
 10" x 1"

Plants

Cordia sebestena
 Euphorbia heterophylla
 (all linear lft.)
 E. mesembrianthemifolia
 Sesuvium portulacastrum
 Melanthera nivea
 Tournefortia gnaphalodes
 Uniola paniculata
 Hymenocallis
 Agave sp.
 Ambrosia hispida
 Candelaria echinata
 Canavalia
~~E.~~ Conyza canadensis
 Agave sisalana
 Ipomoea tuberosa
 I. pes-caprae
 Sporobolus virginicus
 Panicum maximum
 Catharanthus roseus
 Sesuvium portulacastrum

Birds

Mourning Dove

Casuarina for most part.
 Ambrosia hispidula
 forms large patches
 with a few poor Iguana
 plants. Iva on beach
 ridge.

Sporobolus fairly
 general but not
 abundant. Tournefortia
 occasional on sand ridge.
 Suriana is locally
 dead or partly so
 around Ambrosia patches.
 Ipomoea tuba local
 on Suriana.

Near s. end an opening
 in Suriana has a
 mixture of Cyperus, of
 Euph. mesen. Euph. heteroph.
 Boerhaavia, Ipomoea tuba

Nearer end the opening
 is dominated by Uniola.

Along west side is lower
 newer (?) sand flat,
 with sparse Uniola,
 Iva, scattered Suriana,
 etc.

Higher toward end
 being undercut. (photo)

Little veg. except sparse
 Uniola under Casuarina
 (photo)

West half more or
 less open igave sisalans.
 Patch of Panicum maximum.

Beach rock along central
 part, perhaps intertidal
 or just below. One strip
 several beds. Northward
 comes closer to surface
 and near old Cal. cistern
 is above surface of water.

a high sand ridge
 x m. + all along n half
 of w. coast.

Much ~~of~~ *Suriana*
 scrub about 1 m. tall
 much *Agave sisalana*
 much *Opuntia*

The central part and
 around the old Carnegie
 laboratory is very much
 disturbed, no significant
 vegetation pattern. *Casuarina*
 and *Croton* abundant here
Cordia thickets, also, and
Hymenocallis.

There seems to be no
 rock nor gravel above
 high tide level on this
 key. Beach rocks below.
 The sand in general is
Halimeda (sample)

Winds during winter
 said to be from n.w. very
 strong, at times for many
 days at a time. Changes
 outline of keys very much
 at times.

general - Tortugas Keys
 Birds -

On Bush Key are very
 numerous ~~to~~ sooty terns,
 a great many of
 them fully grown young.
 Said to leave in middle
 Sept. and return in April.

Small colony of frigate
 birds on Bush Key.

A white heron with yellow
 bill on Garden Key, flew to
 Bush Key.

A willet on Long Key.

A large number of common
 noddies on Bush Key, nesting
 with mostly almost
 grown young, fully feathered.

Several turnstones
 along beach, Bush Key.

Several Royal terns
 flying bet. Garden
 and Loggerhead.

Aug. 3 - Loggerhead Key
~~Flat low sand inlet~~
 on calcareous sand

- 43026 *Salvia*
 1 around lighthouse building
 1 27 *Bursera simarouba* Sarg.
 occasional in Suriana scrub
 1 28 *Capriaria biflora*
 rare around lighthouse buildings
 3 29 *Sporobolus virginicus* L.
 common generally
 1 30 *Euphorbia prostrata* Ait.
 around lighthouse building
 1 31 *Euphorbia hirta* L.
 around lighthouse building
 1 32 *Eragrostis amabilis* (L.)
 local around lighthouse building
 4 33 *Cyperus planifolius?* L. C. Rich.
 common in openings in
 Suriana scrub.
 5 34 *Solanum*
 very local in Suriana scrub
 3 35 *Boerhavia*
 occasional in openings
 3 36 *Usnea*
 common on dead Suriana twig
 3 37 (lichen)
 common on dead Suriana twig
 1 38 *Uniola paniculata*
 dominant in many openings
 2 39 *Ambrosia hispida* Pursh
 dominant in certain openings
 4 40 *Iva imbricata* Walt.
 locally common on sand
 ridges and flats

flowers blue-purple.

shrub 2 m. tall, sterile.

forming loose sod in sand

prostrate, purplish.

prostrate

small tufts.

shrub 2 m. tall; ripe
 fruit bright red.

prostrate; flowers
 rose-purple.

bright green

culms erect

prostrate, inflorescences
 erect.

— dense shrub to 0.6 m. tall,
 leaves bright green.

- 43041 *Cenchrus*
3 very local in opening in turiana
- 1 42 *Caesalpinia*
very local in turiana scrub
- 3 43 *Tournefortia gnaphalodes*
common locally
- 3 44 *Ipomoea pes-caprae* (L.) Roth
occasional in openings
- 2 45 *Cenchrus*
occasional in openings
- 3 46 *Euphorbia heterophylla*
common in openings
- 1 47 *Panicum maximum*
local, one patch near buildings
- 2 48 *Ficus*
one tree near lighthouse
- 1 49 *Argemone mexicana*
one plant near lighthouse
- 2 50 *Hymenocallis*
very common
- 3 51 *Chloris petraea* Sw.
very local, near buildings
- 3 52 *Ipomoea tuba* (L.) G. Don
occasional in turiana scrub and in opening
- 3 53 *Scaevola plumieri* (L.) L.
local, 2 patches and a few bushes on beach and sand flat.

- culms procumbent,
tough
tangled vine; sterile.
- (L.) R. Br. low bush
- prostrate; flowers
bright purple.
tufts of spreading culms.
- much branched herb,
leaves variable, bracteal
leaves red at base,
up to 1.7 m. tall.
- large tree, fertile branchlets
on trunk.
- branched herb.
- somewhat caespitose,
~~scapes~~ leaves distichous,
scapes terminal, sharp
edged, perianth white,
~~stamens~~ stamens webbed.
- twiner; flowers white,
faded in late forenoon.
- shrub 0.6 m. tall; leaves
fleshy; flowers white.

Garden Key - on top
of wall

Boerhaavia

Cenchrus echinatus

Cenchrus

Bidens leucantha

Stachytarpheta

Sida procumbens

Conyza canadensis

Capriaria

Cynodon

Melanthera

Euphorbia mesem.

Eragrostis domingensis

Ipuntia diirenis

Portulaca oleracea (red form)

Ipomoea pes-caprae

Casuarina (seedling)

Chloris petraea

Canevalia rosea

Bursera sumatrana (seedling)

Cyperus planifolius

(from p. 25)

All of keys west of Key West except a few bars and one small one e of Boca Grande are completely covered by mangroves.

Key West is completely covered by city, but many little mangrove patches in water north of city. Extensive weedy land around as still.

East of Boca Chico a great complex of keys. On north side hundreds of little mangrove patches, locally forming a maze, some larger, all mangrove-covered. Tidal channels in them mostly open at both ends. Eastward the mangrove keys have more exposed bar flats around them.

West of Flamingo Lable Is. is cut by a number of canals, dug many years ago in an attempt to drain the swamps. These seem to cause mud to be stirred up in the waters they connect with.

Aug. 3 - Garden Key
on flat sand on north
side of Key

- 43054
4
1 55
2 56
4 57
1 58
~~4~~
1 59
3 60
3 61
2 62
3 63
2 64
1 65
1 66
1 67
- Cyperus planifolius* R.Br. L.C.
common
- Cyperus planifolius* R.Br. L.C.
occasional
- Chloris petraea* Sw.
very local
- Canavalia*
occasional
- Avicennia germinans* (L.) L.
seedling established at top of beach
- same -ⁿ top of wall of Fort Jefferson
- Cenchrus echinatus* L.
occasional
- Cenchrus*
occasional
- Boerhavia*
common
- same - inside Ruined Fort Jefferson
- Euphorbia*
sheltered spot
- planted tree
- planted tree
- Lawsonia inermis*
planted tree
- Heliotropium curassavicum* L.
very local
- Desmodium canum* (Brael.) Schinz + Thellung
local

- Rich. caespitose
- Rich. culms ascending to erect.
vines, flowers magenta.
- culms depressed, spreading.
- culms depressed.
- stems prostrate to
slightly ascending,
becoming paniculate
- erect, tips arching.
- small tree, 6 m. tall
- small tree, 4 m. tall.
- small tree
- glaucous; flowers white.

40

1963 Florida

- 43068 *Waltheria indica* L.
rare
- 69 *Psidium guajava* L.
planted shrub

Aug. 5 - south of Homestead
along limestone road
embankment above ditch.

- 70
rare
- 71 *Kosteletzkya virginica*
locally common at water edge
- 72 *Eupatorium coelestinum*
common

Aug. 5 - Key Largo, 8 miles north
of Lake Surprise

- 73 *Trema lamarckiana*
common in roadside scrub
on raised reef limestone.

Aug. 5 - Key Largo, 6 miles north
of Lake Surprise.
in low forest on raised reef limestone

- 74 *Callicarpa*
occasional
- 75 *Mourneria ovata*
occasional
- 76 *Santana involucrata*
occasional
- 77 *Morinda royae*
common
- 78 *Quettarda*
common

to p. 46

Dry Tortugas atoll 41

erect; sterile.

shrub 2 m. tall; flowers
white.

rays white

~~flowers~~ erect, main stem
unbranched; flowers pink.
heads purple; herbage
not aromatic.

shrub 0.5-1 m. tall.

erect, sparsely branched,
ripe fruit purple.

small tree, flowers white,
fruit orange, fleshy.

shrub 1.5 m. tall;
fruit purple.

shrub, to 1 m. tall;
flower white, fruit turning yellow.

tall shrub; fruit
immature.

Aug. 5 - trip from Miami to Key Largo
Mangrove road across
everglades separates the
influence of salt water, which
results from drainage, from saw grass.

12625

on oolitic limestone
fill along road across
mangrove swamp
Spermatoc
Euphorbia
Eustonia scalatum

12633

Jewfish Creek.
Lake Surprise
Key Largo - emerged reef.
Scrub forest up to 5-6 m.
tall, dense but irregular
in places. Much *Metopium*.
Bursera, *Piscidia*
Pulcherrima abundant.
Chiososy, *Psychotria*
Mouinda, *Jurettarda*

Trees about 1-2 m
apart 5-8 cm dbh. Irregular
sparse shrub layer
1-2 m tall. Nothing on
ground except sprouts
and seedlings, not many
of these.

Vines - *Vitis rotundifolia*
Chococosa, fairly abundant

ground surface broken
coral rock, with a
considerable accumulation
of litter and partially
decomposed litter, giving
a false impression of
smoothness on surface
(Photos - begin 62-10 perhaps 8 frames)

Opuntia
Bullonia
Borreria
fruticosa

Beyond this, after an
almost imperceptible
change to lower elevation
the forest is replaced
by a *Palbergia* scrub
about 1 m tall, with
scattered emergent trees.
Then a zone of *Batis*
with scattered shrubs,
this only about 20 m wide.
Then a strip of mangrove,
Rhizophora, *Laguncularia*
Avicennia, about 20-30 m
wide, ^{5-6 m tall} ~~much taller~~, then
the sea.
(Photos of edge along road).

photos

Farther north, on the
higher ground, the forest
becomes taller, *Lysiterna*
dominant & at least common.
On the bay side the forest
is larger, 8-10 m; this gives
way to mangroves.

The reef has a cap of a dense soft limestone, smooth on the surface. Forms rather thin patches at about the edge of the Batis and somewhat into the *Dalbergia* scrub. Coral shows through locally. (samples).

On east side of road north of Key Largo large area of dwarf mangrove about 1 m. tall (*Rhizophora*). Do not grow taller except on disturbed soil of road embankment, where they reach several times as tall. The scrub formed by the *Rhizophora* is closed or almost so. No obvious explanation of dwarfing.

- 43079 *Dalbergia*
2 abundant, forming a scrub in open places, undergrowth in forest
- 3 80 *Chiosoclea alba*
abundant in undergrowth
- 1 81 *Quettarda*
occasional
- 3 82 *Psychotria*
common in undergrowth
- 1 83 *Callicarpa*
occasional

Aug. 5 - 8 mi. n. of Jewfish Cr.
on coral fill road bank
in mangrove swamp

- 1 84 *Eustonia exaltata* (L.) G. Don
rare
- 1 85 *Spermocarp*
common
- 3 86 *Euphorbia blodgettii*
common

scrambling shrub

vine

shrub; fruit immature.

shrub 0.8 m. tall

unbranched, flowers whitish

very glaucous, flowers
light purple.
erect, flowers white.

Labels written

Aug. 17 - Sleepy Hollow

in ^{tall} bottom-land forest

43087 *Tipularia discolor*
occasional

88 *Aplectrum hyemale*
rare

89 (grass) *Cinna arundinacea* L.
common locally in more open spots

90 *Cryptotaenia canadensis*
occasional

^{several}
pseudobulbs pale grayish,
scape erect, leaves none
at this season; flowers
dull brownish purple, column
pale green.

pseudobulb single, pale
grayish; leafless at this
season; scapes erect, fruit
pendent.

^{culms}
erect, solitary.

erect

1962 District of Columbia

Aug. 14 - Kenilworth Gardens

Large series of ponds of varying sizes from a few yards across to half an acre or so, with various species of *Nymphaea*. *Nelumbo* in some - in one or two a very handsome species of *Jussiaea* with a large bright yellow flower, a slender fruit, and stems densely soft pubescent. *Hibiscus moscheutos* and other shrubs around edges. Many of the *Hibiscus* are large pinks or even red varieties. Deep red ones very handsome.

Pis pseudocornu also common in margins. A purplish-green, dirty looking bloom is very obvious on some of the ponds.

Between the ponds and the Anacostia River is a strip of ^{tidal} swamp, with *Salix nigra*, *Ulmus*,

Cornus serotina, *Fraxinus*, *Ilex rubrum*, *Betula nigra* - different ones dominant in different areas.

Zizania aquatica in margins, also *Impatiens*, *Rambunct*, *amaranthus*

N. lotus
+ *N. lutea*
+ hybrids

Viburnum
much
Vitis

tidal
Around the south and southeast side is a broad strip of apparently natural marsh more or less dominated by *Typha*, but with abundant *Polygonum punctatum* and *Impatiens capensis*, some *Amaranthus cannabinus*, a little *Zizania*, and another large grass (at least superficially like 43089), and *Sagittaria* scattered in the marsh. There are patches of *Alnus serrulata*, *Hibiscus moscheutos*, *Cornus amomum*, either mixed or in separate patches. Along the path, a strip of more or less open water with *Peltandra virginica* and ~~*Sagittaria patens*~~ some *Pontederia*.

The *Typha* marsh has the vegetation in 2 strata - the *Typha* about 2 m tall, the *Zizania* and *Amaranthus* about the same; the *Polygonum* and *Impatiens* and *Sagittaria* forming a definite layer about half as tall. Locally *Zizania* is dominant. Locally *Rhynchospora*.

1962 Virginia
Aug. 19 - Orkney Springs

Mixed forest, pine
oaks, etc. with a
low shrub layer of
Gaylussacia baccata,
Vaccinium stamineum,
etc. mostly Ericaceae,
some small *Nyssa*.

The forest is very
dry. Moss very
abundant, forming
in places large patches
or an almost continuous
layer, this dry, where
made of *Leucobryum* it
is cracked into segments.

Aug. 20 - upper part of
slope from Big Meadows
to Fisher Gap.

open to closed
second growth low
scrub forest - of
Betula populifolia,
Pinus spp. *Prunus serotina*,
Quercus spp. *Fraxinus*,
Liriodendron, *Pyrus*,
etc.

Many herbs in
flower, esp. in openings.

Chrysanthemum leucanthemum

Allium cernuum

Solidago arguta

Monarda clinopodia

Aquilegia canadensis

Daucus carota

Prunella vulgaris

Silene stellata

Aster macrophyllus

Achillea millefolium

Oxalis (europaea) ?

Centaurea maculosa

Hedysotis purpurea v. *tennifolia*

Plantago lanceolata

Impatiens pallida

Eupatorium rugosum

Arctium minus

Polygonum persicaria

Oxianthus armeria

Lobelia inflata

Solidago bicolor

Cimicifuga racemosa

Carolina
atriplicifolia
medeola
virginica

54

1962 Virginia

Aug. 19 - 1/2 mile ^{west} south
of Old Rag View Overlook,
on Skyline Drive. ^{near route 47}
in spring choked ^{with}
with sedges and moss.

30
42391

Chelone

x2

x3 92 Scirpus
common

x1 93 Carex
common

x1 94 Habenaria
common

x1 95 Epilobium
occasional

x1 96 Botrychium
rare

x1 97 Hydrocotyle americana L.
common

x1 98 Juncus
common

Aug. 20 Big Meadows,
slope toward Fisher Gap
in low scrubby second
growth woods.

x2 99 Viola palmata L.
common in dense shade

43100 Galium circaeans Michx.
rare

x2 01 Luzula
very local in moist densely
grassy place on slope

Shenandoah Nat. Park 55

same place where
Spiranthes was abundant
in spring.

large clump, stems
about 1 m. long; corollae
white with dull purple
orifices.
root crown deeply buried.

part flowering - root
fleshy, horizontal.
flowers pink.

flowers ^{light} red.

base of stem and roots orange
culms solitary.

Labels typed 1962 Virginia

56
by N.K.
43102
71

Solidago arguta Ait.
common

Aug. 20 - slope west of
Big Meadows, Trail to
Lewis Falls

in mixed oak forest

03 *Thelypteris hexagonoptera*
locally common

04 *Thelypteris*
rare

05 *Coralorhiza maculata*
rare and local

06 *Monotropa hypopitys* var.
rare

07 *Liparis liliifolia*
occasional

08 *Monotropa hypopitys* var. *americana*
occasional, locally common

09 *Goodyera pubescens*
very local, under oaks

Aug 20 - low hill north
of Big Meadows swamp
in low mixed second-
growth woods.

10 *Amphicarpa*
locally abundant in
openings

11 *Lycopodium complanatum* var. *tristachyum?*
local

ctd. on N. 60

Shenandoah Nat. Park 57

erect, flowers bright
yellow. (petiole winged)

stem bronze, flowers
bronze, lip white with
deep purple spots.
stems reddish.

capsules winged.

stem ^{part} yellow, flowers
dull yellow.

leaves firm, gray-green,
veins white; flowers white.

tangled in herbaceous
vegetation; affected
by powdery mildew; small
tubers on rhizomes; flowers
dull rose purple.

forming a close mat,
rhizomes apparently varying
from buried to almost superficial.

Aug. 20 - Big Meadows,
slope west from
trailer camp and picnic ground
Thick second growth
woods of oaks, Betula
allegheniensis, Juglans nigra
Prunus serotina, etc.

Plants in flower (not ~~noted~~ ^{noted} above)

- Laportea canadensis
- Hieracium paniculatum
- Campanula divaricata
- Goodyera pubescens
- Monotropa uniflora
- Monotropa hypopitys.

Woods ^{and meadows} around swamp
plants in flower (not noted
above)

- Hieracium scabrum
- Aster macrophyllus
- Solidago puberula or nemoralis
- Rhus glabra
- Lactuca (yellow)
- Cirsium discolor

In swamp - in flower

- Cirsium odoratum
- Sanguinaria canadensis
- Cicuta
- Spiraea latifolia
- Vernonia glauca
- Lobelia cardinalis
- Epilobium
- Hypericum mutilum
- Rudbeckia laciniata
- Eupatorium perfoliatum
- Conium maculatum

- 43112 *Solidago*
occasional in openings
- 13 *Scirpus*
occasional (commoner in swamp)
- 14 *Isoetes*
common in dried pools
- 15 *Licuta?*
common (abundant in openings in swamp)
- 16 *Thalictrum discolor* L.
common (also in swamp)
- 17 *Conium maculatum?*
occasional (also in openings in swamp)
- 18 *Juncus*
occasional ~~in~~
- 19 *Epilobium*
occasional
Aug. 20 in drainage ditch at upper end of ^{Big meadows} swamp in wet sandy mud
- 20 *Epilobium*
common
- 21 grass
abundant
- 22 *Eleocharis*
common
- 23 *Juncus*
occasional
- 24 *Juncus*
occasional

stream

flowers yellow.

caespitose

flowers white

erect.

erect,

caespitose, fruits maroon

flowers pink

decumbent at base.
 small tufts, culms erect,
 dense tufts
 small tufts
 dense tufts

1962 Virginia

43125 *Gratiola neglecta*
common

Aug. 20 lower end of Big
Meadows swamp
in Sphagnum
matted in other vegetation

X2 26 *Habenaria*
locally common

X1 27 *Lycopus*
common

X2 28 *Galium tinctorium*
common

X2 29 *Epilobium*
common

Aug. 20 west slope of
Swift Run Gap

in flat valley bottom
in deep woods.

X1 30 *Epipagus virginianus*
uncommon - under large *Fagus*
grandifolia

X1 31 *Coralorhiza*
rare

X7 32 *Orchis spectabilis* V.
occasional on ~~dry~~ dry ground

X4 33 *Hydrocotyle americana* L.
locally abundant in mud
along stream

X1 34 *Lobelia puberula* var. *simulans* Fern
rare

stems very fleshy,
prostrate; flowers
white

rhizomes fusiform,
stems filiform; flowers white,
erect; weak; flowers
white, both 3- and 4-pid.
flowers pink.

plant bronze, buds purplish.

plant bronze, buds unopened;
base bulbous, above the
white coral-like rhizome.

forming a tangled mat

flowers blue-
purple

1962 Virginia

43134a *Ranunculus*

x2

rare in wet ground along stream

x1 35 *Saxifraga micranthidifolia* (H. Arn.) Britt.

locally common on wet ground along stream

x3 36 *Mimulus alatus* Ait.

locally common on wet stream banks

x2 37 *Galium lanceolatum*

very local on dry rocky ground

x1 38 *Geum canadense*

occasional

Aug. 20 - Pinnacles

Picnic Ground

x5 39 *Carya ovalis* (Wang.) Long.

occasional

Shen. Nat. Park

erect

inflorescence dry

flowers lavender

stems almost erect.

petals white.

small tree 3 m. tall;
fruit immature; with
prominent beels in
upper half.

Aug. 23 - flight from Minneapolis to Portland
 Land west of Minneapolis is generally green, has many lakes. They vary from clear and dark to muddy. The land is largely cut into a patchwork of mostly rectangular fields, blocked off into squares by section-line roads. These fields are mostly cultivated. There are some small patches of wood and wooded stream bottom lands and lake margins. Many patches of wood and (probably) marsh reflect old ponds and stream courses. Some fields are tawny yellow, others gray, others black. Some lakes are partly changed to bog. No large patches of forest.

Westward the amount of forest rapidly diminishes. Perhaps the amount of uncultivated land - pasture not cut into small patches increases. Still many lakes.

In a short time trees are almost lacking. Poor visibility to about the Dakota border. Then a landscape (from 39000') with myriads of small ponds or potholes. Westward these diminish rapidly in number, until they are very sparsely distributed or absent.

Very soon the fields assume an almost completely rectangular pattern, though not uniform in size or orientation. Almost all n-s-e-w, but largely long and narrow, more e-w alignment than n-s. More brown than green.

Poor visibility

From time to time through openings in the clouds the landscape is seen to change from a completely cultivated one to more and more prairie with only stream ravines as discernible features.

East of the Black Hills white erosion scars are prominent. The Black Hills show up as an almost completely black wooded area.

West of the Black Hills the terrain, though at first very flat, shows more and more relief. No creeks until we cross a river that conspicuously meanders through a wide, somewhat wooded, partly cultivated bottom-land. Otherwise no cultivation, or locally large patches of dry land farming, indicated by rectangular pattern lighter than the general brown of the landscape.

As the relief becomes more pronounced narrow gallery forest appears along streams, then considerable dark forest in areas of mountains. Flat plains between mountain patches are grassy. Occasional cultivated, green areas in river bottom lands.

Generally, the rougher, more rocky the land the more wooded, except for some steep eroded slopes and cliffs. These

generally south and west facing. North and east slopes generally wooded.

Westward, flat topped mesas and buttes become prominent features.

Then really rough mountains - prominently dendritic drainage patterns flowing generally north. Relation between these and distribution of forest not clear or obvious.

Main roads show up, curiously, in red, even in a dominantly white or light gray and tawny landscape.

Forest greatly diminishes westward.

Broad strips of green cultivated land along north-south trending river.

West of this mesa-land with black, wooded complex stream drainage going east from it and west on and from it.

Another green stream valley, joining the east one far to the north. It runs s.w. + n.e. south

of it dry, treeless, grassy high plains - gray-brown.

Far to north some forest. Stream now and now from west with wide cultivated area.

Westward vast plains with some relief locally. Dry cultivation on flatter areas. Mostly dry grass. More and more cultivation westward - yellow grain fields, but curiously banded - dark and yellow in subequal strips.

Far to north mostly forested.

Along course and for a distance northward the same striped pattern in large patches, forming a mosaic with dry grassland. The darker stripes in the cultivated area appear to be similar dry grassland.

No more forest to north.

Some narrow gallery in ravines forming a w-e. dendritic pattern.

Billings
mont.

Then bare ravines still some striped cultivation.

Then treeless brown plains and small mountain ranges, narrow green cultivated bottom lands.

Then grassy plains with prominent w-e. drainage, gallery along ravines.

Then a tremendous mountain range mostly wooded, but with a large sharp peak that is bare, with snow on its north summit slopes.

West of this prominently cultivated land predominates, but with much dry grassland too, the grass thin and landscape eroded.

Cultivated patches become more and more sparsely scattered.

Far to north some prominent but low, strike scarps.

Then a densely wooded mountain mass.

Then much yellow

grains, fading out to dry grass on country with some relief. This more accentuated westward, to a prominent n.e.-s.w. valley and ridge strike-slip system.

Winding between two sections of this a river, with a wide rather green basin directly on course with a small city and an intricate system of streams meanders - probably marshy or formerly so.

Then broad grassland, high plains with scattered grain fields and some relief.

A n.-s. narrow green bottom-land with a tightly meandering stream. Then a high but rather narrow mountain range about half wooded. A smooth grassy valley, then another largely wooded mountain range. This one very extensive

and with only minor areas of grass.

Butte Montana at west base of these mts. Range open but mine. High plains west of Butte. Country to north wooded and rough.

They more wooded mountains, then a ~~rough~~ wooded range - the ~~Bitterroot Mts.~~

Northward there become bare and eroded.

Then the tremendous Bitterroot Range, mostly densely wooded except some very large peaks which evidently extend above the forest, and are grassy or bare.

This vast forest extends a long distance, but far to north is grassland.

Then mostly forest, as far north as can be seen.

Road systems in parts of the mountains very intricate - no apparent reason.

Then mountains become locally bare and grassy over large areas, forest, where wooded, becomes sparser, ^{valley} then an

1962 9 dabo

area of tremendous ruggedness, high peaks with patches of snow, deep east-west canyons. Forest sparse on roughest part, dense on more moderate slopes to westward.

Then forest becomes very sparse and steep mountains are largely bare. Forest still on gentler slopes. Much forest to north.

Country now a very rough jumble of mountains and a mosaic of bare rock and forest of varying density. Steeper slopes tend to be bare.

Westward of dense forest becomes more continuous.

Some local mountain parkland. Road scars on forested slopes very conspicuous.

Relief becomes gentler but forest almost continuous and ~~very~~ dense for a long distance.

Then more open country but still predominantly

forested.

To northward a large open but rather rough area - flatter parts in grain, becoming dominant westward. Forest still along course, but gradually fading out to grassland.

Canyons still with north facing slopes wooded.

Country along course still mountainous, to north a large rather level basin, mostly cultivated.

This bounded on s.w. by Snake River Canyon - a tremendous bare gash in the land. South of it, and locally north of it large patches of forest.

West and south very rough mountains, practically treeless, the Snake R. canyon meandering through them. Northward a forested area north of that the cult. basin continues.

Then along course, south of Snake and

partially wooded mountains; north of them and west of Snake, high bare mountains continue. North of them broad plain, cut by canyons, level parts in grain.

Along course the landscape now is a high plateau but very deeply dissected, leaving flat mesas on which are grain fields. Some forest on moderate slopes. Steep slopes bare and brown. Spectacular deep canyons cut this. Plateau becomes densely wooded.

To north the dissected plain continues, mostly in grain. Between it and forested plateau, very rough partly wooded mountains. These then extend south at least to course. become less continuously wooded. Forest in canyon bottoms and on north slopes.

Then forest mostly disappears. The

Oregon - Washington 77
belt of dissected plain swings south to course.

Columbia River Basin. High Peaks of Cascades in distance to west (200 mi.) landscape now about $\frac{1}{2}$ to $\frac{2}{3}$ in grain, rest bare and brown except some bottom lands of a westward flowing river (Snake?)

Texture of country along course interesting - main streams flowing w. n. w. to Columbia. Tributaries flowing s. w. and n. e.

Columbia here is a great reservoir, lower bottom lands of tributaries broad, green, cultivated.

Dam across Columbia when it flows west, just above a tributary from south.

Away from river and tributaries vast plains, brown, with some yellow grain fields locally. Some rough, eroded areas. Strip of grain well to north of Columbia R.

Mt. Rainier, Mt. Adams,
~~Mt. St. Helens~~, ~~Mt. Jefferson~~
 on west horizon.

Vast brown plains
 continue along both
 sides of Columbia -
 large areas of grain
 away from river.

In distance to north
 and well ahead a large
 forested area, with
 grass north and south
 of it. Forest continuous west to Cascade.

Columbia comes closer
 to course, even below
 the dam a very impressive
 river. As it approaches
 course it becomes
 incised, with prominent
 cliffs. Grain fields
 and grasslands (fallow
 land?) form a vast
 mosaic. This extends
 along the terraces
 along both sides of
 river.

A large island in river.
 Here river is again very
 wide, obviously backed
 up by a dam.

Some lumber on south side.

Course crosses river.

Country on north terrace
 mostly grassland.

River swings north
 again. Terraces become
 dissected and partly
 wooded.

The Dalles.

Mt. Hood on left, also
 Mt. Jefferson.

The ^{light} mountains are
 all pine cones.

Country generally wooded,
 rough, some scrub
 patches. Cleared right-of-way.

Patches of fine forest -
 tall conifers, some spruce,
 some bare peaks and
 rock ridges, old
 fire scars, logged
 off areas, - latter
 especially extensive
 n.e. of Portland, - in fact
 almost continuous
 over large areas. More
 slashed off land, by
 far, than forest in
 the region north of
 Columbia and east
 of Portland.

Approaching river
 land flattens out
 and forms a mosaic
 of cultivation, pasture,
 some forest in canyons.

Large grayish marsh
 just north of river.

Aug. 24 - trip from Corvallis to coast, by auto.
west from Corvallis (rt. 20)
Philomath - cultivated land with some large oaks.

West of this a range of hills partly wooded with mixed second growth. A common oak that looks like *Q. lobata* in valleys, with *Alnus*, *Carex macrophyllum*, *Fraxinus* sp., *Conifers* on slopes and hilltops.

Higher up, most of forest is mainly coniferous. On the western slope of the range, near Burnt Woods a mosaic of lower broadleaf wood with conifers.

Toledo Salt Marsh -

Expanse of tidal marsh, perhaps 30 acres.
Largely a matrix of grass, with admixture of *Atriplex patula*, and locally *Salicornia*, patches of *Gnaphalium*, *Potentilla anserina* etc.
Some large rotting logs provide a substratum for *Gaultheria shallon*, *Pseudotsuga* ~~seed~~ saplings and stunted trees, *Achillea*, *Vaccinium*, *Polypodium*.

Deep brown muddy soil.

Yaquina Bay State Park Bluffs with *Pinus contorta*, low trees up to 4 m. a few small *Pseudotsuga*. Thick undergrowth at least locally of *Gaultheria shallon* with some *Smilax involucrata*.

Newport

Meadow of *Carex abrupta* and *Potentilla anserina*, up to 2 m. tall in wetter spots, back of active dunes and in front of

high stabilized dunes, these with *Rubus*, *Alnus*, *Gaultheria*, *Sonchera* var. *Picea*, *Pinus contorta*, etc. Grassy meadows also, with clumps of *Pinus*. *Pteridium*

On active dunes - *Cakile*, *Abronia*, *Armeria*, *Convolvulus soldanella*, *Polygonum paronychia*, *Fragaria chilensis*, *Lathyrus*, *Vicia*, *Potentilla*, *Solidago*, etc.

Cakile one of most pioneer species.

Photos of *Pinus contorta* and of general landscape showing wind shear. Also of *Carex* meadow.

Outside of dunes a broad sand flat.

Wreckage all over this flat and back in dunes.

1 mi. n. of Waldport

A number of parallel rows of high dunes, upper slopes mostly active, locally wooded, many species of psammophytes on active parts. Forest, mostly *Picea*, in depressions and on one outer ridge, this being badly sand-blasted.

In forest in depressions, also on unwooded steep sides, a dense scrub or undergrowth of principally *Gaultheria*, up to 3 or even more m. tall.

Many photos - ending rolls of Kodachrome (K-6) and B. & W. (62-10).

Aug. 24 - Toledo, ~~Salt Marsh~~
near mouth of Yaquina River

in tidal marsh

43140

X²

Grindelia stricta
common

X²

41

Halicornia ambigua
common along tidal channels

X²

42

Halicornia ambigua
common along tidal channels

X²

43

Pyrus fusca Nutt.
occasional at edge of

X¹

44

Juncus
common

X¹

45

Galium aparine L.
common in edges of

X¹

46

Salix
common around

~~common to salt marsh~~

X¹

47

Glauis maritima
common locally

X¹

48

Faurea carnosus
rare

X¹

49

Orthocarpus castillejoides
occasional

X¹

50

Triglochin maritima
common

X²

51

Pseudotsuga menziesii
in edge of

X¹

52

Aster
common

stem woody at base,
sending up herbaceous
branches; rays and disk yellow.
stems terete, fleshy; fls.
staminate

stems terete, fleshy;
flowers pistillate

small tree, fruit
more or less cylindrical,
immature.

stems slender, terete.

stems weak, reclining
on other plants; fls. white.

shrub 3 m. tall; sterile.

sub-fleshy

fleshy; flowers yellow

subfleshy; bracts white-tipped

leaves subterete; spike erect.

~~shrub~~ shrub 3 m. tall,
spreading; leaves white beneath,
stems erect from
horizontal rhizomes,
rays lavender, disk yellow,
turning reddish.

- 43153 *Polypodium vulgare* v. *occidentalis*
 on rotting log
 X₂
 X₁ 54 *Vaccinium watsonii*
 on rotting log
 X₁ 55 *Gaultheria shallon*
 common on rotting logs

Aug. 24 - about 2 miles
 south of Yaquina Bay Bridge

sand dunes

- X₂ 56 *Rhododendron macrophyllum*
 common in low woods on fixed
 X₂ 57 *Carex obnupta*
 dominant in moist
 meadow between
 X₂ 58 *Fragaria chiloensis*
 common on
 X₂ 59 *Franseria bipinnatifida* Nutt.
 locally common on
 X₂ 60 *Arctostaphylos uva-ursi*
 common on
 X₂ 61 *Pentstemon ramosissimus*
 common on
 X₃ 62 *Armeria maritima*
 common on
 X₂ 63 (grass)
 local on
 X₇ 64 *Franseria bipinnatifida* Nutt.
 locally common on
 X₂ 65 *Calcile edentula* (Bigel.) Hook.
 common on outer, moving

fronds very young.

small shrub; ripe
 fruit black.

small shrubs; ripe
 fruits black.

shrub 2.5 m. tall.

erect, to 2 m. tall.

runners extensive

prostrate, forming
 mats, inflorescence
 ascending.

prostrate, forming
 mats

prostrate, forming
 small prickly mats
 flowers bright pink

prostrate, forming loose
 mats, leaves fleshy.
 fleshy, much branched
 herb; fruits terete, swollen
~~flowers~~ light purple
 petals

1962 Oregon

- 43166 *Polygonum paronychia*
common
- X1 67 *Calsila maritima* Scop.
rare, on outermost
- X1 68 *Rumex maritimus* var. *frugosus* (Phil.) Duran.
rare, on outermost
- X2 69 *Goodierya oblongifolia*
local ~~in~~ in open pine
woods on fixed dunes
- X2 70 *Maianthemum bifolium* (L.) ^{var. *hammichaticum* (Opunt.) Pehr.}
common in pine woods
on fixed dunes

Aug. 24 - 1 mile north of
Waldport Lincoln Co.
on high dunes of fine sand

- X2 71 *Anaphalis margaritacea*
locally abundant
- X2 72 *Agoseris aparigioides* (Less.) Green
occasional
- X1 73 *Poa macrantha*
common
- X2 74 *Glehnia leiocarpa*
common
- X3 75 *Carex macrocephala*
abundant, binding sand
- X2 76 *Franseria chamissonis* Less.
common
- X2 77 *Abronia latifolia*
common
- X1 78 (grass)
common in sheltered ^{open} places

flowers pinkish white

leaves fleshy; petals
purple; fruit terete, not
conspicuously swollen.leaves with irregular
white along midrib;
flowers white.fruit fleshy, finely
red-dotted.erect stems from
buried rhizomes; involucre
papery white.

caespitose

stolons very extensive

fleshy; flowers white

rhizomes deeply buried

prostrate below, distally
ascending, forming
loose mats.prostrate, forming extensive
loose mats, flowers bright yellow.

- 43179
42179 ✓ 2 Polypodium vulgare var. occidentale
local on steep sand slope on
- ✓ 2 80 Salix
common in depressions between
- ✓ 2 81 Poa macrantha
common on
- ✓ 2 82 Pteridium aquilinum var.
common on ~~top~~ active

Aug. 25 - 3 mi. n. of Florence
at junction with Mercer Lake Rd.
in Thuja plicata swamp

- ✓ 2 83 Blechnum spicant
abundant
- ✓ 1 84 Ledum columbianum
common in undergrowth
- ✓ 2 85 Darlingtonia californica
abundant in water and
common on Sphagnum hummocks

same - in pine woods
- ✓ 3 86 Myrica californica
common in undergrowth
- ✓ 3 87 Arctostaphylos columbiana
common in openings

Aug. 25 - Cape Perpetua State Park
in dense mossy ^{some} forest

- ✓ 2 88 Selaginella oregona
abundant but very local on
tree trunks.

shrub 3 m. tall; sterile.

caespitose, stems scarce or
lacking.

rhizome deeply buried,
fronds all about this size
(farther inland, on fixed
dunes much larger, more
conspicuously hairy beneath)

sterile fronds spreading,
young ones without
anthocyanin; fertile fronds erect.
shrub 2 m. tall, sparsely
branched; flowers white.
leaves and scapes erect;
top of leaves with translucent
spots.

shrub 2.5 m tall, ~~aromatic~~ ^{fragrant};
fruit green.

shrub 2 m. tall, bark
smooth, dark red brown.

pendent

Aug. 25 - Cape Perpetua Viewpoint
 Large promontory with steep bare cliffs and irregular forest of Picea on top and upper slopes. On south side aspect is rather dry - close undergrowth of Gaultheria.

Cape Perpetua State Park - canyon with big spruce forest - luxuriant undergrowth largely of Gaultheria, but ferns very abundant - Blechnum, Polystichum, Athyrium cf. filix-femina, Dryopteris dilatata (on log). Selaginella oregona very abundant on a leaning tree trunk, pendent. Mosses and liverworts abundant on trees, also lichens.

South of here the coastal bluffs are covered by dense matted low scrub of Gaultheria. Some areas of low spruce forest, badly damaged by wind and spray.

Southward the slopes are grassy. Then scrub and low forest of Pinus contorta with some spruce. This matted and windsheared on exposed areas. On steep slopes the scrub is mostly Picea.

South of Sea Lion Caves Alex covers bluffs.

Inland somewhat is a tall forest, to 30 m. of Pinus contorta, ~~Libocedrus~~ ^{Thuja}, etc.

Undergrowth of Gaultheria, Rhododendron, Vaccinium, Arctostaphylos.

Low swamp of Thuja and a few Pinus contorta - open and mostly not more than 5-8 m. tall. Undergrowth of Ledum, Myrica, Tsuga, Gaultheria, and abundant Darlingtonia in wet Sphagnum and in water. On hummocks Blechnum.

Honeyman State Park -

Enormous active dunes, separated from a strip of smaller dunes ~~to~~ back of

the beach. The large dunes are gradually moving into the forest - largely *Pinus contorta* about 20 m. tall. Some spruce. *Alnus* and *Salix* are common. Heavy undergrowth of *Gaultheria*, *Vaccinium*, *Myrica*.

Much taller forest inland - 40 m. mostly spruce, hemlock, *Thuja*. with tall understory of *Myrica*, *Rhododendron*, *Vaccinium* 3-4 m. tall.

— Inland from Florence along st. 26. up Suislaw River. Hills along sides of river, covered with coniferous forest, but considerable part has been logged off and is covered by scrubby *Alnus* forest. *Acer macrophylla* is common with *Alnus* on lower slopes. Valley bottom in pasture, where not flooded by dams. *Alnus* common in fence-rows and on river banks.

Farther up the valley considerable areas on slopes have been logged clean

and are now replaced by a chaparral-like scrub.

Just west of Triangle Lake 40 m. e. of Florence

Partly logged *Pseudotsuga*, with some *Thuja*, *Alnus*, etc. Undergrowth of *Vaccinium parvifolium*, *Acer circinnatum*, *Corylus*, etc.

Polystichum muricatum abundant on ground, not much else in denser parts of forest. Some *Pteridium*.

Interior valley - flat cultivated land, scattered *Quercus garryana*, patches of wood - *Quercus*, *Fraxinus*, etc. Grassy rolling hills with scattered oaks and patches of oaks; small bushes

Aug. 20. Silver Falls State Park
Beautiful vertical drop.

Pseudotsuga - abies forest, 50 m.
Understory of *Acer macrophyllum*,
A. circinnatum, *Alnus*.

Aug. 25 - just south of
Triangle Lake
in Pseudotsuga forest

- 43189 *Montia sibirica*
in clearing
- ✓ 1 90 *Prunus*
edge of clearing
- ✓ 2 91 *Alnus oregonia* Nutt.
abundant in edges of
- ✓ 2 92 *Galium triflorum*
common
- ✓ 1 93 *Asarum*
very local in paths
- ✓ 2 94 *Pinnaea brealis*
local on steep dry bank
- ✓ 2 95 *Polystichum munitum*
abundant on ground
- ✓ 1 96 *Vaccinium parvifolium*
common in undergrowth
- ✓ 2 97 *Acer circinatum*
common in undergrowth
- ✓ 3 98 *Corylus cornuta* var. *californica* (A. DC.) Hook
occasional in undergrowth
- ✓ 2 99 *Vaccinium parvifolium*
common in undergrowth
- 43200 *Athyrium filix-femina*
14301 occasional, terrestrial in
moist place

flowers white, ^{turning pink} petals emarginate

small sterile tree, 6 m. tall,
bark rather silvery gray.

small tree, leaves
glaucous beneath.
procumbent

rhizome prostrate, under
~~leaf~~ litter, with strong
ginger odor when crushed.
prostrate

fronds spreading to
ascending
slender shrub 2 m.
tall, branchlets in one
plane; fruits ^{dark} coral red
shrub 3 m. tall, stems
slender, green.

sterile shrub 1.5 m tall.

shrubs 2 m tall;
fruits coral red.

1962 Oregon

Aug 31 - trip Corvallis - Arcata
via Eugene, Redport

Freeway east of Corvallis
and south goes through
extremely flat land,
intensively cultivated -
small patches and
strips of forest of
Fraxinus regina with
some *Quercus garryana*
along sloughs and in
areas of poor drainage.

Where the fields are
left in grass there is
an invasion of shrubs -
possibly *Populus* *pusilla*
or *P. trichocarpa*. Willows
along ditches.

Southward forest
becomes much more
abundant, much of
it young and thickly.
Hills are scattered over
plain and are more or
less wooded with conifers.
The thicket on the flats
is *Fraxinus*, *Quercus*,
etc.

South of Eugene
the flat Willamette
valley pinches out
and the hills get close

together. Low Douglas fir
on flats as well as on hills.
Acer ~~floridum~~ macrophyllum
becomes common along
with *Quercus*, *Fraxinus*,
Salix, etc. in thickets
and forest.

Road across mountain
to coast along canyon of
Elk Creek - Douglas fir
second growth with
much admixture of
Acer, etc. Patches of
grass and brush on
steep slope.

Rhus diversiloba forms
an irregular understory
in disturbed or ^{open} areas -
is a shrub 3-4 m tall,
robust but climbing when
it is in contact with trees.

Below Elkton hills, where
denuded of trees, *Cytisus*
scoparius forms a scrub,
but only locally. Many
such areas are grassy
with *Pteridium* on talus
slopes, not where rock
is close to surface.

Much of it a scrub
with *Holodiscus*, etc.
And *Quercus garryana*
especially around

Umpqua
River

grassy places on steep rocky bluffs. Here many dead been openings even in pre-logging times, where the oak could have lived. Oak patches also on certain level places at foot of slopes, where there may have been Indian villages. The grass is doubtless very

Farther down the slopes have a mixed forest of maple, alder and Douglas fir, this replaced by thick Douglas fir forest.

In valley bottom alders, etc. have some moss on trunks. Some steep bluffs logged clear.

Much of the area is rather fern forest, pure pseudotsuga.

On flats along river well up sitka spruce pines also Acer circinatum comes in as understory on flat, also up under Douglas fir on slope. Where

river valley broadens out there are remnants of excellent sitka spruce forest.

Estuary extends up 12 or more miles. Tidal influence shows in a practically vegetation-free belt several feet wide on steep banks.

Down just above Reedfoot are marshy flats with Scirpus, Carex, Arthrospora, etc. In this area hemlock (*Tsuga heterophylla*) almost replaces the Pseudotsuga, as an associate of *Picea sitchensis*.

Dune area at Hausee fairly extensive area of active dunes. Patches of *Ammophila*, scattered *Poa maritima*, *Clombe* of *Falis*. Patches of *Arctostaphylos uva-ursi*. Patches of wood - *Pinus contorta* with *Alnus*, esp. in depressions. Some *Arctostaphylos columbiana*. On active parts also *Tanacetum* sp. in patches, scattered *Lupinus arboreus* (mistle) patches of *Rhacomitrium ericoides*.

High dunes burying spruce forest. Some trees dead, others buried to within 3m. of top, but still living, fruiting.

About 10 mi. n. of Coquille - *Chamaecyparis lawsoniana* common in forest.

South of this forest is a thicket of *Alnus* with some emergent conifers.

Extensive salt marshes at Bandon, on s. side of Estuary of Coquille R.

S. of Bandon a large area of *Ulex europaea* 2-3 m. tall - in an old fire. fairly extensive, flat area.

A broad strip, several miles wide, is of fixed dunes along most of coast. Some of them covered by a scrub of *Arctostaphylos*. Some wooded with *Pseudotsuga*, etc. understory of *Rhododendron macrophyllum*.

Southward along coast the bluffs and seaward slopes are grassy. Probably secondary.

South of Pistol River extensive fixed dunes with *Lupinus arboreus* which is dying from disease (or insect attack).

Del Norte Redwood Grove drier type has undergrowth of *Gaultheria*, with *Asarum*. Much *Alnus* + *Rithrocarpha* in understory. Occasional Douglas fir. More moist type has ~~*Clintonia andersonii*~~ *Oxalis oregona*, *Blechnum* ^{*atryanum*} and *Polystichum munitum*. *Clintonia andersonii* and *Scoliopus* are character species. Drier types *Rhododendron macrophyllum*, etc.

In *Oxalis* type much vine maple.

A belt of Sitka spruce between redwood and coast.

Sept. 1 - auto trip, Eureka to Bull Creek

Flood plain of Eel River. Presently only flooded in lower part - up a few miles rejuvenation has gone on so that the river, even in flood is confined to ravine. Was covered by Sitka spruce up to about

then gradually replaced by redwood; also on sides of valley, upward replaced on sides by Douglas fir, tar oak and madrone + oaks.

Eel River meanders from side to side of valley undercutting sides, forming rather bare cliffs of poorly consolidated sediments.

Above Scotch mostly second growth redwood - this after old style logging with bullocks, etc. Modern logging with heavy machinery so destroy the environment that redwood scarcely regenerates.

Some virgin redwood along highway, but this will be largely destroyed by widening of highway.

Polytrichum subopacum (L.) G. & G.

in alluvial flat

Rubus stratosus (L.) W. & A.
Anemone vitifolium (L.) G. & G.
Aconitum deltoideum (L.) G. & G.

Urtica glabella (L.) G. & G.
Galium triflorum (L.) G. & G.

Trifolium arvense (L.) G. & G.
Trifolium repens (L.) G. & G.

Smilacina stellata (L.) G. & G.

Presently badly affected by highway - camping, etc.

first redwood forest - Rockefeller forest and gravel along ~~Eel River~~ ~~Rockefeller~~ next ^{road} - Bull Creek flat grove with old gravel deposit with willow, ~~along~~ ^{along} ~~ss.~~ ^{ss.} ~~of~~ ^{of} ~~Eel~~ ^{Eel} ~~River~~ ^{River}.

next road. 6 beginning of b.w. 67-12 inside Rockefeller grove

flat
Enormous trees, ± 2 m. dbh. 75 m. tall; closely spaced 1-10 m. but canopy not continuous, usually Noctiflyte above basal few m. in redwood.

ground flora -
Oxalis reginae
Pteridium aquilinum (L.) Kuhn
Polystichum muricatum
Woodwardia
Athyrium filix-femina
Ficella unpoliata
Blechnum spicant
Pentaria tenella
Claytonia sibirica
Carex stipata (result of deer damage dist.)

second row
Umbellularia (10 6-9 m)
Acer macrophyllum (mainly along stream)
Much bare ground.

Bull Creek is carrying down great amounts of gravel from 1955 logging, which fills up the creek bed so it cuts laterally, undercutting the trees - perhaps 800 trees lost on Bull Creek in Rockefeller grove

Mounds around the bases of trees, up to 7-12" high, are earth, with only a few cm of litter on top.

Above these flats, ~~the~~ ~~from~~ this far inland, on the slopes the forest changes immediately to a mixture of

area near "Tallest Tree"

Ground flora

Achlys triphylla (red, not named)
Tridentalis
Galium triflorum
Blechnum spicant
Polystichum munitum
Dicranum smithii
Tiarella
Corylus inopercinata
Rithocarpus
Acer

Hard ground with very thin *Coralis*, ^{*Tiarella*} scattered *Polystichum*, patches of *Achlys*,

Trees ^{to} 17 (or 20) feet dbh.

Sept. 1 - air trip Arcata - San Francisco
sand flats opposite
airport show some wind
striation.

Solid fog layer to the
foot of the mountains,
for a short distance
south, to just n. of section.

Tops of coastal hills
tend to be bare, especially
on seaward slopes, but this
clearly result of clearing
in most cases. Because of
angular outlines.

Forest on west slopes sparse
because of logging (many
small roads forming a
network. Some quite bushy.

The bare patches tend
to be stable - little invasion
of woody plants on many
of them, others quite bushy.
Brushy ones show evidence
of recent logging in scattered
sticks lying around.

Course is well inland
with rugged wooded mountains
westward. More and more
open grassy areas southward,
also more recently logged
areas. Tremendous
variation, from dense
forest to dense scrub, to
sparse forest, sparse

scrub, savanna, grass -
with little evidence of
pattern, little relation
to topography. Network
of logging roads and
scars very conspicuous.
A few patches of good
timber still unlogged.
This course follows
S. Fork Eel River.

South of Angelo Place (?)
as a number of very
fine uncut small
drainages, densely wooded,
but badly logged near
one interfingered with them.

Well inland a very
conspicuous strip that is
mostly grass. A large
area seaward of it
is ~~to~~ mostly uncut.

But roads follow ridges
around it, and logging
spreading from it.

Along ~~some~~ principal
stream bottom lands
are strips of redwood, where
they have not been logged.

In these uncut areas
there still seems to be a
rather clear pattern, but
the logging obscures it.

Another large uncut
area a little farther south,
a few prairie patches - not topographic scars.

much of this patch seems to be redwood.

Inland a fine mosaic of grass and forest on rough country.

Then course goes over bare ridges with more bare area on west or south-west slopes. Some brush, some mainly the ravine are wooded. Some cultivation. Ukiah

Wooded ridges to west.

South of Ukiah the wooded ravines on grassy west slopes are apparently oak. Conifers only west of valley in which Ukiah located!

Chaparral, rather thin on many hills, esp. on east and north slope, more grass on west & south. Little continuous forest areas of any size from here on except some steep on e. slopes.

Southward more and denser chaparral. Many burned areas. Locally incredible erosion.

Then suddenly much more forest, but only a mosaic of patches with chaparral and grass.

Inland a broad, rather cleared valley. Course not far from coast. Some restricted areas of redwood.

A meandering rather large river - meander very ancient and deeply entrenched in rough mountains runs toward coast.

Somewhat south of this the forest suddenly stops and dry rolling hills ^{land} with gallery forest in larger ravines fills area east of the large southward running cultivated valley (Napa??) This ~~is~~ rolling land brown with numerous wind breaks, a few small ponds. is very extensive. Southward it becomes rough and many headwater drainage are wooded or scrub covered, tailing down into gallery wood in ravine below. Bay is ~~to~~ south - prob. San Pablo Bay. The large gallery ended at Black Point? Large complicated reservoir just east of course.

South of this larger patches of forest and dense scrub, and more numerous ones, tending to coalesce. Here another large reservoir, very complex. Coniferous forest on west side, chaparral with patches of forest on east. The west side ~~runs into~~ the e. slope of Mt. Tamalpais. W. slope here is chaparral, but w. of Tamalpais is a large area of grass with some oak trees and patches. Ridges tend more and more to be low southward.

Thin chaparral and small patches of wood in ravines on Angel I.

A great tongue of fog extends in from Golden Gate clear across Bay. Fog covers hilly part of S. F. but flat part and bay are clear

Baccharis pilularis

Sept. 3 - Santa Cruz Mts.

The east slope, from Saratoga up to Saratoga Pass is ~~is~~ mostly wooded, with some redwood not far below the summit. The grass is dry but other things are mostly still green - ferns etc. The wood is of live oak (mostly *Quercus agrifolia*), *Acer macrophyllum*, *Quercus kelloggii* (esp. on upper slopes), *Umbellularia*, *Arbutus*, etc. The ~~top~~ summit ridge s.e. of Saratoga Pass to the Fred Hertz ranch, is semi-open, semi-wooded, with *Pseudotsuga*, *Quercus kelloggii*, *Arbutus*, *Acer macrophyllum*, *Baccharis pilularis* etc. *Rhus diversiloba* locally common, both erect (to 2 m. tall) and climbing.

The grassy and herbaceous vegetation of the open areas, rocks and roadside, is rather dry, but such plants as *Madia*, *Zauschneria*, *Chrysopsis*, *Fragaria*, *Potentilla*, *Haplopappus*, *Polygonum*, etc. are still green. *Eriogonum* locally common along road. *Acer* fruiting heavily.

Aug. 31 - Hauser, 4 mi. n. of
Cross Bay, Coos Co.
on extensive active dunes
of fine sand

43201 *Tanacetum*

✓1 common, forming patches

✓4 02 *Salix*

common

✓2 03 *Salix scouleri*

common

✓1 04 *Spiraea douglasii*

rare on shallow ditch

✓3 05 *Polygonum paronychia*

common

✓1 06 (grass)

rare

✗

same - in moist open
low place in active dunes

✓2 07 *Juncus*

common, forming patches.

✓1 08 (comp.)

very local in patch of *Juncus*

✓1 09 *Eleocharis*

rare in patch of *Juncus*

✓2 10 *Juncus*

common, forming small patches

✓2 11 *Sesyrinchium*

very local, in patch of *Juncus*

✓1 12 *Aster*

very local, in patch of *Juncus*

✓1 13 *Bidens cf. frondosa*

local in patches of *Juncus*

spreading by underground
stems; flowers yellow.

densely branched shrub

1 m. tall.

densely branched shrub

2 m. tall.

unbranched shrub 1 m. tall.

flowers pink.

flowers pink

rhizome deeply buried,
culms cylindric, wing;
not erect, emerging in
lines from sand.

culms erect from buried
rhizomes.

flowers bright yellow

ray flowers purple, disk yellow
turning red.

erect, disk flowers yellow

- 43214 *Agrostis*
rare, in patch of juncus
- ✓ 15 *Ranunculus*
rare, in patch of juncus
- ✓ 16 *Hypericum* cf. *mutilum*
occasional in patch of juncus
- ✓ 17 *Veronica*
common

Aug. 31 - about 10 mi. north of Coquille

- 3 18 *Chamaecyparis lawsoniana*
common in coniferous forest

Sept. 3 - Hayward, Alameda Co. Calif.

- 1 19 *Amaranthus*
common in hard ^{dry soil} ground and full sun along street

Sept. 3 - summit ridge of Santa Cruz Mts., Summit Road, 4 miles s.e. of Saratoga Gap.

on dry roadside in chaparral with *Pseudotsuga* and *Quercus kelloggii*

- 3 20 *Haplophragma arboreum*
occasional in road cuts
- 3 21 *Eremocarpus setigerus*
locally common in full sun
- 1 22 *Acer macrophyllum*
common
- 2 23 *Zauschneria californica*
local
- 1 24 *Polygonum*
rare, in full sun

creeping, rooting at nodes, ~~no~~ flowers bright yellow. petals yellow, plant tending to be reddish.

flowers purplish.

tree 20 m. tall, branches drooping & pendant.

prostrate

2900'

Santa Clara - Santa Cruz County boundary

aromatic shrub 1 m. tall, flowers yellow, rays none.

small tree, 6-8 m. tall

large clump, flowers scarlet.

stems ascending

118

1967 California

43225

Heterotheca

1 common in full sun

1 26 *Corylus cornuta* var. *californica*
occasional in edge of thicket

1 27 *Madia*
in open grassy place

3 28 *Madia*
common in open grassy flat place

1 29 *Madia*
in open grassy steep slope

1 30 *Madia* cf. *madisoides*
on steep rocky bank

3 31 *Solidago*
common on roadsides
and other open places

~~37~~

Santa Clara - Santa Cruz Co.

119

small clumps, ^{usually erect} of a few stems;
flowers yellow
shrub 2 m tall

rays yellow

plant strong-smelling;
ray flowers yellow.
ray flowers yellow

rays yellow

erect, plant gray-green,
flowers yellow.

Sept. 4 - flight from S.F. to Chicago
in Boeing 707.

Low lying cloud layer
over Bay area, extends inward
past the first hills east of
the Bay.

These are brown with
blackish forest in ravines
and steep slopes.

Area north of Suisun Bay
is of brown rolling hills
with no trees except tiny
planted clumps.

Sacramento Valley area a patchwork
of green and brown cultivation,
but eastward large area
e. of Sacramento of brown rolling land
with green cultivated
valley bottom lands
along rivers.

Foothills more and
more wooded with
spruce to locally dense
wood, probably oak
woodland. Flatter
areas mostly grassy
and light brown or
cultivated. This sloping
flat surface more and
more dissected eastward
and more completely
wooded. Remnants of
the old surface grassy
but increasingly scarce

upward. Higher cleared
areas showing conspicuous
contour farming, some of them
green at higher elevations.

Higher areas are light gray
rounded n-s ridges only
very sparsely wooded, almost
completely bare and almost
white ~~east~~ west of Lake
Tahoe. Immediate slopes
around lake are forested.
Lake an intense blue except
a few narrow shallow
areas which are light green.

Forest belt east of lake
dense but narrow, only
on first several ridges.
Then everything is brown.
Then a cloud ~~belt~~ layer.

Then Nevada Basin and
range topography with
bare brown mountains,
rusty to white dry lakes,
reservoirs, a few green
bottom lands and culti-
vated areas. Area around
Fallon cultivated, ~~and~~ a
green and yellow patchwork.
Carson has a pale gray
smooth extensive dry lake
bed with a few darker gray
patches and a tiny patch
of low hills near east side,
some sand dunes there and

a row of them along east side at foot of high range of mountains. Higher parts of these are rather sparsely wooded. Another smaller dry lake east of this range, and vast alluvial fans and washes. Mt. ranges east of this wooded on higher slopes, sparsely so and lower ranges only slightly. Some dry farming east of these in wide valleys, but only scattered enormous fields. A few ~~few~~ smaller bright green fields.

Increasing cloudiness. Mountains mostly too low to be more than brown and very sparsely if at all wooded. Vast alluvial fans, small dry lakes, very locally ^{on fans} much inexplicable road construction forming close network. Elsewhere only a few roads.

Mountains mostly with rounded, not too rough ridges. Locally some sharper erosion. This with scattered trees.

Cloudy.

Then rougher broad valleys, with stream courses green, with some woody vegetation.

Elko Nev.

Then a very high, very rough range, with sharp ridges, some wood, pale gray rocks, snow patches in sheltered high spots.

Much cloudiness.

Then a high range to south, quite densely wooded at higher elevations. Then broad alluvial slope down to a very extensive almost white featureless area - Bonneville Salt Flats, or Salt Lake Desert.

This is enormous. Some small patches and strips of dunes eastward. These varying in character and pattern. Some are partially fixed - but by what? Some small patches of darker bare, mountains like islands. Then some obviously fixed tiny dunes.

Then a reddish alluvial slope and a range of mountain with vegetation on lower slopes, upper ridges bare.

Small outlying ridges in alluvial slope to east.

Then another wide desert valley, but with flow patterns rather than a dry lake. Some green patches of cultivation to south.

Then a higher range of mountains, wooded in higher parts, to south broad alluvial slopes, and a small area of cultivation patchwork at south end of Great Salt Lake ~~to~~, this lake bounded by salt flats and, in one place, evaporation basins.

Another mountain range and an extensive cultivated plain, with salt lake city.

Then extensive area of thinly wooded mountains possibly mostly chaparral-covered, but from 37000' this is hard to be sure of. This mountain system is very complex, with complicated drainage and ridge patterns.

A valley with bottom lands (cult.) and a reservoir perpendicular to course.

More complicated mts.

with widely spreading drainage basin draining westward into valley just mentioned. Quite extensive, irregularly vegetated, but thinly where relief is little, densely on higher areas. Cloudiness.

Desert drainage systems, incised in broad alluvial slopes, between low drainage desert mountains, ^{the remaining} turning eastward, but eventually entering a n-s. river.

East of here low mountains, often more or less flat-topped, trending n-s, fading gradually into almost a plain with low n-s. scarps, and flat topped ridges, and ~~at~~ ^{beyond} Rock-springs, Wyoming, only rather minor relief. All desert, but with a conspicuous pattern of spots of vegetation.

Eastward this changes to low desert mountains, densely wooded mountains visible to south. ~~Here~~ locally plain, locally low mountains. Then

higher mts, locally
densely and extensively
wooded, especially
to south, much grass
a desert on more gentle
relief along course,
only patches of grass
to south, locally
closely imbricated
e-w strike ridges.

Then drainage to
n.e. over vast alluvial
slope, dissected into
low mesas near
mts. Low ridges
running east or n.e.,
scattered ponds with
whitish (alkaline?) margins.
Country all brown,
treeless, sparsely
hilly in a complex
pattern, changing
to a plain with a few
rather sizeable lakes
(a reservoir).

Then abruptly, a low
mountain range, and
east of it, a somewhat
dissected plateau, and
more low mountains
interspersed with valleys
with a little relief.
Drainage east or north of east.
Hills lower and lower,
high plains, still cultivation, poor visibility.

Interfluvies are low mesas
with complex, incised
dendritic drainage pattern.
Large areas of cultivated land
on lower terraces.
Edges of mesas conspicuously
whitish.

East of Scotts Bluff, a
great multitude of ponds
or reservoirs and some
dune areas in grassland.

Sept. 8 - near Williamsburg
headw. of Powhatan Cr.

Swamp with *Acer rubrum* and *Nyssa sylvatica*. Shallow wide ravines cut into the coastal plain, total relief perhaps 8-10 mi. with many spring heads.

Rich herbaceous flora, many ferns, rich shrub layer.

Slopes with *Liquidambar*, *Quercus prinus*, *Pinus taeda*, *Acer rubrum*, *Q. velutina?*, *Fagus*, etc. Some of pines and tulips are quite large.

Sept. 8 - headwaters of Powhatan Creek, near Williamsburg,

in swamp

- 43232 *Rudbeckia laciniata* var. *digitata*
✓ 4 wet ground, dense shade
- ✓ 33 *Elephantopus carolinianus*
wet sandy roadside
- ✓ 2 34 *Cyperus retrofractus* (L.) Torr.
wet sandy roadside
- ✓ 1 35 *Desmodium*
wet ground, dense shade

get county from map James City Co

rays yellow; disk greenish yellow

flowers lavender

loose clumps with single ascending culms. erect

1962 Virginia - Ohio - Md.

Sept. 16 - Sleepy Hollow

Fauquier Co.

weedy vacant lot.

48236

Eupatorium

local, occasional

label
writtenSept. 22 Dryast Woods,
2 mi. W. Centerville

Belmont Co. Ohio

✓1

37

*Galium*occasional in shade
of tall mixed hardwood
forestSept. 30 - Arnold, just north
of Annapolis, Anne Arundel Co.
in mixed second-growth
hardwood forest, mostly *Liriodendron*

✓12

38

Asplenium platyneuron (L.) Oakes

very local, on earth banks

✓1

39

Botrychium dissectum

on low flat ground

✓4

40

Myrica cerifera

common in edges, along roadside

✓1

41

Botrychium virginianum

rare on low flat ground

✓5

42

local on roadside

slightly gregarious
because of rhizomatous
habit; erect; heads white.

branching at base.

population sample,
all plants seen.shrub 2 m. tall,
fruit lead-gray.prostrate twining
vine stems several m.
long; fruit terete.

Cathedral State Park

126 acres

hemlock (on 21' girth)
 red maple
 white oak
 yellow birch
 sugar maple
 black cherry

seven acres on west
 side (27500 trees (ft.))
 on gentle n. slope -
 bordered on west by
 meadow. Border of
 thick small growth
 of hemlock, red oak, red
 maple, birch, yellow
 birch, tulip

thick undergrowth
 of small trees opens
 in about 50' or more
 then first large hemlock
 and undergrowth
 immediately opens up
 but with scattered
 suppressed trees - large
 trees well spaced but
 canopy fairly complete

ground cover of patches
 of Dryopteris spinulosa,
 Renontactis, Rypogonium,
 Luccidium, Lilla, Mitchella,

Large trees
 hemlock dominant
 red maple
 black cherry

A large red maple with
 conspicuously twisted trunk.

Cherry 3' dbh

fallen chestnut.

Several small areas of
 wind damage. Hemlocks
 have been broken and windthrown.
 thick growth coming up here.

Sept. 22 - Dysart Woods

broad ravine with old mixed hardwood forest - fairly open beneath, canopy essentially complete trees to 30-35 m. tall, varying in size to ~~20~~ 1.8 m dbh. mostly much smaller, down to 10-15 cm. for canopy trees, a few smaller suppressed ones.

Many seedlings & saplings up to 2 m tall. Almost no herb layer

Quercus alba
Acer rubrum var. *glabrum*
Fagus grandifolia
Acer rubrum
Nyssa sylvatica
Prunus serotina
Fraxinus americana
Liriodendron
Alnus sp.

Some beeches have small branches along trunk.
 Bark with numerous tubercles 1-2 cm wide, 1 cm high on bark, arranged

Herbs

Botrychium virginianum
Sedum ternatum
Epipagus
 a *Aster macrophyllum*
Aster sp.
Prenanthes trifoliata
Solidago caesia
Polygonatum pubescens
Galium trifolium
Galium lanceolatum

Cornus florida
Ilex toxicodendron
Parthenocissus

Sept. 29 - Suitland Bog
 Access difficult - the old way all built up. Coming from the back the steep slopes are covered by dense *Kalmia* and *Lonicera*, hard to penetrate.

Sept. 30 - Falls Church area
trees are just beginning
to change color - red
maples are, some of them,
beginning to show
red at ends of branches.
Platanus somewhat
brownish or yellowish.
Nyssa, Cornus, Prunus
serotina have turned
or are turning red. Most
Liquidambar still green.
Liriodendron in driest
situations show some
yellow.

Sept. 30 - Head of Broad Creek,
Anne Arundel Co. Md.

Nyssa in full color,
abundant around shore
at head of creek. Father
O'Neill says that in normal
years the leaves would
be gone by now.

Crabs abundant in
2-3' water, esp. on sand or silt
bottom. About 50 males or
1 female.

One blue heron, several
gulls.

Sept. 30 - Arnold, north
of Annapolis.

Second growth hardwood,
very ragged and irregular,
with scattered pines.

The prominent understory
of Cornus is turning red.
Nyssa almost completely
turned. Liquidambar
beginning to turn purple.
^{some fruit purple.}
Sassafras beginning
to turn red. Tulip with
some yellow leaves, ^{some trees} almost yellow.
Platanus brownish green.

Red maple beginning to turn.
Carpinus beginning to turn.

The interesting thing
is that almost all species
vary so much in how
far they have changed color
from tree to tree.

This woods largely
Liriodendron. On very
low rolling hills and
flat low ground.

Soil brown sandy loam
with some fine gravel
and iron concretions.

Botrychium on low
flat ground.

Much honeysuckle
locally. Still flowering slightly.

Oct. 7, Rattlesnake Point Overlook
Mixed deciduous forest
Fraxinus, Carya, Ostrya
Quercus rubra, Prunus, etc.
Juglans nigra, Robinia, Tilia
Shrub layer of Hamamelis,
Crataegus, (edggs).

Hamamelis flowering
abundantly, Clematis
virginiana fruiting.

Fraxinus, Carya, Prunus
serotina well colored.

Quercus rubra, Robinia,
Tilia turning.

Oct. 7 - Rattlesnake Point Overlook,
Skyline Drive (Northern section)

43243

✓ 3

Crataegus chrysoarpa Ashe
local on talus below road
greenstone

tree-like shrub 3 m. tall,
fruit bright red, fleshy,
rather tart.

Oct. 10 - Washington, D.C.

✓ 2

44

Crataegus
planted around park

small tree 3 m. tall;
fruit scarlet.

19 200

Mr. Hugh Mathison

owns

Castello Hammock

Homestead

15 local species of ferns.

should also try to get

"Lincoln Hammock" next

to Castello.

Two together form the

most interesting hammock

in Fla.

Mathison interested in

conservation, but afraid

people are taking advantage.

